

Chapuis-Lardy, L., Vanderhoeven, S., Dassonville, N., Koutika, L.-S., Meerts, P., 2006. Effect of the exotic invasive plant *Solidago gigantea* on soil phosphorus status. *Biology and Fertility of Soils* 42, 481-489.

Accepted version

Last version before publication

Effect of the exotic invasive plant *Solidago gigantea* on soil phosphorus status

Chapuis-Lardy L.^{1,2,*}, Vanderhoeven S.¹, Dassonville N.¹, Koutika L.-S.¹, Meerts P.¹

¹ Laboratory of Plant Genetics and Ecology, Université Libre de Bruxelles, B-1160 Brussels, Belgium; ² Current address: IRD, Institute for Research and Development, BP 64501, F-34394 Montpellier, France; * Author for correspondence (e-mail: lydie.lardy@ird.fr; phone: +33-467416228; fax: +33-467416294).

Abstract

Invasions by exotic plant species can modify biogeochemical cycles and soil properties. We tested whether invasion by early goldenrod (*Solidago gigantea*, Asteraceae) modifies soil phosphorus pools at three sites in Belgium. Aboveground phytomass and soil samples (0-10 cm) were collected in early goldenrod patches and in adjacent, uninvaded, grassland vegetation. Soil P fractions varied between the three sites in line with corresponding differences in organic matter, carbonate and clay contents. In addition to site-specific impacts, plots invaded by goldenrods generally had higher concentrations of labile P (i.e. resin-extractable inorganic P, bicarbonate-extractable inorganic and organic P). Soil CO₂ release, alkaline and acid phosphomonoesterase (PME) activities were also higher in invaded plots,

suggesting that the increase in labile Pi was due to enhanced mineralization. Phosphorus uptake by vegetation was 1.7-2.1 times higher in invaded plots, mostly due to the higher annual yield of *Solidago gigantea*. Altogether, the results indicate that *S. gigantea* enhances P turnover rates in invaded ecosystems.

Keywords Biological invasion, Early goldenrod, Phosphomonoesterase activity, Sequential phosphorus fractionation, Soil phosphorus availability

Introduction

Over the past two decades invasions by exotic plants have been recognized as a serious cause of biodiversity loss and natural habitat degradation (Vitousek et al. 1997; Levine et al. 2003). In addition to their impacts on biodiversity and community structure, invasions by exotic plants might also modify ecosystem functioning in a more subtle way. The possibility for plants to modify soil properties and biogeochemical cycles is well recognized. Plant-induced heterogeneity in soil properties has been recognised in many types of ecosystems (Hobbie 1992; Van Breemen and Finzi 1998; Ehrenfeld 2003) and the structure of soil microbial communities has been shown to differ in soils beneath different species of plants (e.g. Westover et al. 1997; Kourtev et al. 2002). The functional capacity of the soil microbial community, as reflected in the activities of enzymes involved in nutrient mineralization processes, also varied among soils under the canopy of different plant species (Kourtev et al. 2002; Chen et al. 2004). Indirect effects of plants on microbial processes can equal or exceed direct effects of plant uptake on nutrient retention (Hooper and Vitousek 1998). Vitousek (1990) first pointed out that, like other plants, exotic species could modify soil processes.

To our knowledge, only two studies have examined the effects of exotic plants that are invasive in Europe (Plichta et al. 1997; Mitchell et al. 1997). Recently, Ehrenfeld (2003) and Levine et al. (2003) reviewed 79 and 30 papers respectively, dealing with effects of invasive exotic plant on soil nutrient cycling processes. Most published studies have focused mainly on C and N cycling, whereas effects on P were only rarely examined (Ehrenfeld 2003). Only quantitative changes in total or exchangeable P were reported and no data exist on the impacts of exotic plants on P dynamics. Phosphorus is essential for the basic processes in plants that sustain growth and development. Sequential P fractionation method (Hedley et al 1982 modified by Tiessen and Moir 1993) has been and still are widely used to characterise soil inorganic (Pi) and organic (Po) P forms, based on their extractability in specific reagents, which could be related to their availability and chemical forms. Inorganic P removed by anion exchange resin and 0.5 M sodium bicarbonate constitute a highly available Pi pool in soil (Hedley et al 1982). Dilute NaOH extracts more slowly available Pi associated with Fe- and Al-compounds. The weakly alkaline bicarbonate extraction removes easily hydrolysable organic compounds such as ribonucleic acids and glycerophosphate (Bowman and Cole 1978) while the more strongly alkaline NaOH solution extracts less labile Po associated with humic and fulvic acids. Dilute acid extracts Pi from apatite or octocalcium phosphate (Frossard et al 1995). Hot concentrated acid extracts more stable pools of Pi and Po, some of which may come from particulate organic matter (POM) and may therefore be potentially bioavailable (Tiessen and Moir 1993). The fractionation residue is unlikely to contain anything but highly recalcitrant P. Soil organic P represents between 20 and 80% of total P in the surface layers of soils (Dalal 1977) and it also contributes to the P nutrition of plants primarily after being mineralized into inorganic P. The rate of P mineralization depends on microbial activity and on the activity of microorganisms- and plant-synthesized enzymes (Dalal 1977). Among the phosphatases, phosphomonoesterases (PME) form an important group of enzymes catalyzing

the hydrolysis of organic P monoesters (i.e., mainly inositol phosphates that may account for up to 60% total soil Po; Halstead and McKercher, 1975) to orthophosphate, potentially available to both microbes and plants. Plant roots induce several chemical processes that may modify the mobility and bioavailability of P ions in the rhizosphere, namely: root-induced depletion/accumulation of P ions, acidification/alkalinisation and the exudation of organic acids/anions (Hinsinger 2001).

Solidago gigantea (early goldenrod), a native of northern America, was introduced into Western Europe as an ornamental plant and the first observations of naturalization were reported in the middle of the 19th century (Wagenitz 1964). *S. gigantea* considerably increased its range during the 20th century and has become one of the most serious invaders in Europe (Jakobs et al 2004). In the present study, we examined its impacts on soil P pools and bioavailability in three sites with contrasting soil properties in Belgium.

Materials and methods

The invasive species

Solidago gigantea (Asteraceae) is a rhizomatous perennial forb of up to 2.5 m in height. Its shoots are erect and unbranched except within the inflorescence. Within established stands, population growth occurs almost exclusively vegetatively (rhizomes) as has been shown for the closely related *S. altissima* (Meyer and Schmid 1999). Due to this clonal growth the species can form dense stands that exclude almost all other species. In Europe, goldenrod species are not attacked by herbivores that could control population growth and density and their growth performance and competitive ability were apparently better than in their native range (Jakobs et al. 2004).

Site selection

Three sites were selected (Table 1) in the central part of Belgium based on the following criteria: 1) a well established (i.e. at least 10 year old) and still increasing population of the target species, that formed at least 5 m² monospecific patches clearly distinct from the surrounding vegetation, 2) adjacent (i.e. only few meters apart) zones of uninvaded vegetation consisting of herbaceous native species, 3) a soil auger investigation that suggested a relatively homogenous soil cover in the whole area (i.e. no apparent topsoil difference between invaded stands and uninvaded zones). According to the FAO classification, soils were eutric or calcic regosols with anthropic origin (FAO/UNESCO 1990).

Vegetation and soil sampling

Within each site, six 1-m² plots were randomly located within stands of the target species and six 1-m² plots selected between invaded patches in closely adjacent, uninvaded herbaceous vegetation. Aboveground vegetation was harvested at the peak of biomass production in August. The material (all organs pooled) was oven-dried at 60°C for 72 hours, weighed and finely ground (< 150 µm) prior to analysis.

A bulk soil sample, consisting of 6 cores (4-cm diameter, 0-10 cm depth soil borer), was also collected in each 1-m² plot. The twelve composite bulk samples per site were air-dried and sieved to 2 mm before analysis. Subsamples were dried at 105°C to constant weight to express chemical characteristics on a dry soil weight basis. Another subsample was finely ground (< 150 µm) prior to determination of total C and N. Selected chemical and physical properties of the topsoils are summarized in Table 2. Hereafter, the term ‘Solidago soil’ refers to soils in invaded situations, i.e. under the canopy of *Solidago gigantea*, whereas ‘control soil’ refers to soil of the corresponding, uninvaded situations.

Plant and soil chemical analyses

Dried plant tissue was prepared for P analysis by calcination at 550 °C for 5 h, and digesting the resulting ash in 1 *N* HCl. Phosphorus concentrations were determined on digests by ICP-OES. Results were expressed as total P concentration (mg P g⁻¹ plant dry matter (DM)) and converted into P uptake per unit area (mg P m⁻²) using the plant aboveground biomass data.

Soils were analyzed for total C and total N using a Carbo Erba model NC2100 elemental analyser. Soil P was fractionated sequentially with three replicates per treatment following the modified method of Hedley et al. (1982), as described in Tiessen and Moir (1993), with HCO₃-saturated resin strips (BDH #55164-2S, 9 x 62 mm), followed by 0.5 *M* NaHCO₃ (referred to as Bicarb-P), 0.1 *M* NaOH (OH-P), 0.5 *M* cold HCl (acid-P), and concentrated hot HCl at 80°C for 10 min (hcHCl-P). The shaking period for each of the first four steps was 16 h at 20°C followed by centrifugation for 15 min at approximately 10,000 *g* and followed by filtration. The procedure involved 0.5 grams of soil extracted in centrifuge tubes using 30 ml of each extractant. Methods used for the regeneration of the anion exchange strips, the removal of organic matter from the extracts for inorganic P determination (by flocculation following acidification with H₂SO₄) and the determination of total P (Pt) in solution by oxidation with ammonium persulfate were as described by Tiessen and Moir (1993). The residue containing the most chemically stable P forms (Residual-P) was digested using concentrated H₂SO₄ + H₂O₂. Phosphorus in extracts and digests was determined colorimetrically using the ascorbic acid/molybdenum blue method (John 1970). Po in appropriate extracts was calculated by subtracting Pi from Pt (Po=Pt-Pi). Labile Pi refers to available Pi pools, i.e. removed by resin strips and extracted by sodium bicarbonate (resin-Pi + bicarb-Pi).

Soil CO₂ release and phosphomonoesterase (PME) activities

Ten grams of soil were moistened with 2 ml of deionised water and kept for 7 days. Thereafter the soil was placed in 150 ml sealed plastic jars along with 10 ml of 0.1 M NaOH in a small plastic container. The jars were incubated at 28°C and carbon dioxide (CO₂) trapped in NaOH was measured by titration with 0.1 M H₂SO₄ on the Radiometer TIM 900 Titration Manager pH meter using a ABU 901 Autoburette after 7 and 14 days.

Phosphomonoesterase activity was measured in soil solution buffered at pH 6.5 (acid PME, EC 3.1.3.2) and pH 11 (alkaline PME, EC 3.1.3.1) with disodium *p*-nitrophenylphosphate tetrahydrate as enzyme substrate (Tabatabai 1982). Soil samples were incubated for 1 h at 37°C, the reaction terminated by alkalisation and the absorbance determined spectrophotometrically at 410 nm. Controls were processed on each sample to determine the non-enzymatic hydrolysis of the substrate and to correct for background coloration. Enzymatic activities are expressed as the concentration of the hydrolysis product (µg *p*-nitrophenol (NP) g⁻¹ soil h⁻¹). These results were converted to µmol NP g⁻¹ soil h⁻¹ for easier comparison with literature data (using the molecular weight of *p*-nitrophenol, 139.11 g mol⁻¹).

Data analysis

All data are presented as means of six plant samples or six composite bulk soil samples per situation (i.e. invaded vs. uninvaded within a site). Standard deviations are shown in parentheses or as bars. Randomly selected soil samples were analyzed in triplicate and the analytical error shown to be relatively small (mean CVs of 2 % for total P, 7.8 % for P fractions and 8.5 % for PME activities, respectively). All statistical analyses were performed using the *Statistica* software package (©Statsoft Inc. 2003). Data were log-transformed when necessary to improve homogeneity of variances and the significance level was $P < 0.05$ unless

otherwise stated. Invaded and uninvaded plots were compared by means of paired *t*-tests. Comparisons among sites were performed by means of one-way analysis of variance (ANOVA) and *post hoc* (LSD) tests. A two-way ANOVA was also performed with “site” and “plant cover” (invaded vs. uninvaded) as the fixed main effects for selected soil parameters. Pearson correlation coefficients and principal components analysis (PCA) were used to assess relationships between selected soil variables.

Results

Biomass production and P content of aboveground vegetation

Aboveground phytomass of *Solidago* patches, ranging from 760 to 917 g DM m⁻², was significantly higher than that of the native vegetation (from 252 to 438 g DM m⁻² depending on species composition of the uninvaded situations at the three sites; Tables 1 and 3). There was no significant variation of *Solidago* phytomass among sites (ANOVA; $F(2) = 2.0$; $P = 0.167$). It should be noted that the aboveground biomass of the native vegetation (forb rich grassland) did not vary significantly among sites either (ANOVA; $F(2) = 2.8$; $P = 0.095$) in spite of contrasting floristic composition (Table 1). Phosphorus concentrations in the aboveground components of *Solidago* were significantly lower than those of the native vegetation in Sg1 and Sg2 sites whereas no change was observed for Sg3 situations (Table 3). For all sites, *Solidago* significantly took up 1.7-2.1 times more P per unit area (1211-1713 mg m⁻²) than native vegetation (574-878 mg m⁻²; Table 3).

Soil chemical and physical characteristics of the different sites

There was a marked contrast between soils from the different study sites (Table 2). Sg2 soil was slightly acidic whereas Sg1 and Sg 3 were neutral (pH \approx 7.1) and contained calcium

carbonate (0.4 % in Sg1 soil and 1.4 % in Sg3). Due to their high exchangeable calcium content, Sg1 and Sg3 soils presented a higher CEC (average values between control and *Solidago* soils: 44.1 cmol kg⁻¹ for Sg1 and 46.6 cmol kg⁻¹ for Sg3, respectively) than topsoil of Sg2 (8.3 cmol kg⁻¹) (ANOVA; $F(2,28) = 34.2$; $P < 0.001$). Carbon concentrations ranged from 1.48 to 9.36 % (Table 2) with highly significant differences between soils: Sg2 < Sg1 < Sg3 (ANOVA; $F(2,30) = 175.3$; $P < 0.001$). A similar trend was observed for total N concentration with average values of 0.14 % in Sg2, 0.20 % in Sg1, and 0.36 % in Sg3 ($F(2,30) = 89.9$; $P < 0.001$). Clay contents were significantly lower in the Sg2 soil (2.4 % on average) compared to Sg1 (12.6 %) or Sg3 (15.6 %) (ANOVA; $F(2,29) = 32.3$; $P < 0.001$).

Within sites, comparisons of means (t -tests, Table 2) revealed no significant difference between invaded patches and uninvaded plots for clay content, CEC, total C and N contents. At site Sg2, the pH decreased slightly from 6.5 in the topsoil of uninvaded plots to 5.9 under the canopy of *Solidago gigantea* whereas no significant change was observed in Sg1 and 3 (Table 2).

Sequentially extracted P fractions of soil

The sum of all P fractions (Pt(sum)) ranged from 309 to 1067 mg P kg⁻¹ soil (Table 4) with significant differences between sites (ANOVA; $F(2,30) = 102$; $P < 0.001$) as follows: Sg2 < Sg1 < Sg3. Calcium associated phosphates extractable with dilute HCl (acid-Pi) was 7.8 times lower in the non-calcareous soil (Sg2, ≈ 56 mg P kg⁻¹ soil) than in the calcareous Sg1 and Sg3 (417-470 mg P kg⁻¹ soil); This fraction accounted for 16-18 % of Pt(sum) at Sg2, but reached 40 to 60% at Sg1 and Sg3.

Resin P (i.e. highly labile P) is significantly increased under the canopy of *Solidago gigantea* in all sites: 16.1 vs. 25.6 mg P kg⁻¹ soil for site Sg1 ($P = 0.06$), 19.3 vs. 28.1 mg P kg⁻¹ for Sg2 ($P = 0.03$), and 61.2 vs. 91.5 mg P kg⁻¹ for Sg3 ($P = 0.02$) (Table 4). Bicarb-Pi,

another fraction of labile Pi, was significantly higher in Solidago soil of site Sg3 (81 vs. 57.8 mg P kg⁻¹ in control soil; $P < 0.05$; Table 4). In site Sg1, hcHCl-Po significantly increased from 18.4 mg P kg⁻¹ in control soil to 26.9 mg P kg⁻¹ under the canopy of the invasive ($P < 0.05$). The effect of *Solidago gigantea* is also noticeable on the sum of organic P fraction (Po(sum)) for Sg2 (Table 4; $P < 0.05$). The organic fractions extracted by bicarbonate and soda (bicarb-Po and OH-Po) tended to be higher in soil under the cover of *S. gigantea*, but *t*-tests did not show any significant differences for these fractions. When all sites were pooled, two-way analysis of variance showed a highly significant site effect for all P fractions (Table 5). A significant (at least $P < 0.05$) plant effect was found for resin P, bicarb Pi and Po, and hcHCl Po (Table 5). For all of these variables, the invaded soils had higher mean values compared to the control. There was a significant site x plant interaction for bicarb-Pi, indicating contrasting impacts of the invasive species depending on site.

Soil CO₂ release and phosphomonoesterase activities

Soil CO₂ release during wet incubation was significantly higher in soil under Solidago compared to native vegetation at Sg1 (2.8 vs. 2.4 mg CO₂-C g⁻¹ 14 days⁻¹, $t = -2.81$, $P = 0.018$) and Sg2 (2.5 vs. 2.0 mg CO₂-C g⁻¹ 14 days⁻¹, $t = -2.83$, $P = 0.017$), while there was no significant effect at Sg3 site (Table 2). Enzymatic activities ranged from 140 to 503 μg NP g⁻¹ soil h⁻¹ for acid phosphomonoesterase (Ac-PME) and 95 to 641 μg NP g⁻¹ soil h⁻¹ for alkaline phosphomonoesterase (Alk-PME) (Fig. 1). Analysis of variance and *post-hoc* tests showed a highly significant site effect ($P < 0.001$; Table 5). The calcareous soils Sg1 and Sg3 had highest values for both acid and alkaline phosphomonoesterase activities compared to the non-calcareous, less organic Sg2 (Figure 1). Two-way ANOVA revealed a highly significant plant cover effect ($P < 0.001$) on acid-PME activity. The site x plant interaction was also significant ($P < 0.05$), indicating that the impact of *S. gigantea* varied depending on site

(Table 5). Acid phosphomonoesterase activity was significantly higher ($P < 0.05$) in soil under the canopy of the invasive species in sites Sg1 (336 vs. 190 $\mu\text{g NP g}^{-1} \text{ soil h}^{-1}$; $t = -3.53$) and Sg2 (211 vs. 140 $\mu\text{g NP g}^{-1} \text{ soil h}^{-1}$; $t = -3.52$) whereas no difference was observed for the calcareous soil Sg3 (circa 311 $\mu\text{g NP g}^{-1} \text{ soil h}^{-1}$; Fig. 1). In the acidic soil of site Sg2, the alkaline phosphomonoesterase activity of the soil was unchanged in the presence of *S. gigantea*. Calcareous soils in Sg1 and Sg3 presented higher alkaline phosphomonoesterase activity under *S. gigantea* patches than in the control (Sg1: 641 vs. 379 $\mu\text{g NP g}^{-1} \text{ soil h}^{-1}$, $t = -3.93$; Sg3: 623 vs. 477 $\mu\text{g NP g}^{-1} \text{ soil h}^{-1}$, $t = -3.75$; $P < 0.05$; Fig. 1). The two-way ANOVA also revealed highly significant ($P < 0.001$) effects of plant cover and site x plant interaction for alkaline phosphomonoesterase activity of topsoils.

Relationships between soil P fractions and other soil parameters

Resin-Pi was strongly and highly correlated with Bicarb-Pi ($r = 0.97$, $P < 0.001$) and OH-Pi ($r = 0.093$, $P < 0.001$), the latter being also highly correlated to Bicarb-Pi ($r = 0.96$, $P < 0.001$; Table 6). These three forms of inorganic P were highly ($P < 0.001$) but less strongly ($0.71 < r < 0.85$) correlated to soda-extractable organic P (OH-Po) as well as to C and N contents (Table 6). To a lesser extent ($0.44 < r < 0.51$), their relationships with Bicarb-Po, acid and alkaline phosphomonoesterase activities were also significant (Table 6). Acid and alkaline phosphomonoesterase activities were highly correlated with each other ($r = 0.83$, $P < 0.001$) and with OH-Po ($r = 0.74$ and 0.70 , $P < 0.001$, respectively). Soil pH was not correlated with resin Pi, OH-Pi and Bicarb-Po but with Bicarb-Pi ($r = 0.42$, $P < 0.05$), OH-Po ($r = 0.44$, $P < 0.05$) and acid and alkaline phosphomonoesterase activities ($r = 0.45$, $P < 0.01$ and $r = 0.76$, $P < 0.001$, respectively). In principal component analysis, the first component associated with 63.7% of the variance, displayed strong negative loadings (< -0.7) on several variables that could be grouped into three clusters according to their relative position in the plane defined

by the two first principal components (data not shown): i) OH-Po and PME activities; ii) total C and N contents; iii) inorganic P pools (Resin-, Bicarb- and OH-Pi). The second most important component accounted for 19.3% of the total variance and targeted dominantly Bicarb-Po, and secondarily alkaline PME. Soil pH presented similar absolute correlation with PC1 and PC2.

Discussion

“Site” effect on soil P dynamics

We investigated the impacts of the exotic invasive plant *Solidago gigantea* on soil P status at three different sites. The three study sites have contrasting soil properties that are likely to explain the large differences among them in soil P fractions and phosphomonoesterase activities. Sg1 and Sg3 were neutral soils with calcium carbonate and a higher clay and organic matter content compared to Sg2. In the Sg1 and Sg3 soils, the largest pool of P is the CaCO₃-bound, acid-extractable P (Acid-Pi), that may be partly originated from phosphate rocks or basic slag applications in the past. Despite the influence of calcium carbonates, P-sorption was also impacted by other soil properties such as clay and organic matter contents. The significant difference in clay contents between the studied sites (Sg2 < Sg1 = Sg3) may also be responsible for differences in soil P sorbing capacity (Frossard et al. 1995, 2000). Studied soils also differed in their C and N contents; this variability may also be responsible for differences in P fractions as organic matter comprises various variable charge compounds. Moreover, organic matter mineralization produces organic acids that compete with P for sorption sites on soil minerals (López-Hernández et al 1986; Frossard et al. 1995, 2000). Higher values of phosphomonoesterase activities were found for soils with higher organic matter content and CO₂ release rate (Sg1 and Sg3). Such positive correlations have already

been reported in the literature (Bonmati et al. 1991; Tarafdar and Claassen 1988). As expected, acid phosphomonoesterase was predominant in the slightly acidic soils (Sg2) and alkaline phosphomonoesterase was predominant in neutral soils (Sg1 and Sg3); the influence of soil pH on phosphomonoesterase activities was described previously (e.g. Eivazi and Tabatabai 1977; Juma and Tabatabai 1978).

In-site soil conditions

A fundamental problem in invasion studies is the potential for pre-existing differences in soil properties that made certain areas more conducive to invasion. The experimental design does not allow us to exclude with complete certainty the possibility of such pre-existing differences between uninvaded and invaded zones. However, uninvaded plots were located close to the selected invaded patches, being separated only by a few meters. Moreover, preliminary field observations and samples analyses (e.g. texture and cationic exchange capacity) showed that soils of uninvaded and invaded zones did not differ significantly. It should also be noted that the *S. gigantea* patches are still expanding. In this context, the observed differences in P more probably resulted from the contrasting plant cover of the two types of plots, uninvaded vs. invaded. Our failure to detect significant impacts at some sites (e.g. resin-Pi in Sg1) may be due to the relatively low sample size (n=6). Alternatively, significant site x plant interactions may be due to genuinely idiosyncratic impacts of the invasive species. Thus, pH was found to significantly decrease under the canopy of *S. gigantea* in site Sg2 and not in sites Sg1 and Sg3 where calcium carbonate probably enhances soil buffering capacity.

P uptake by vegetation

In our study, *Solidago* showed differences in plant growth. It produced 1.7-3.5 times more aboveground biomass and took up about twice as much P per unit area as the native plants. Yearly P turnover will be consequently greater for *Solidago* than for native vegetation. This may explain the higher proportion of organic P fractions in invaded patches. In line with its higher productivity, *Solidago* has deeper rooting systems compared to native vegetation (pers. obs.; USDA-NRCS, 2004) which may result in uplift of mineral nutrients from deeper soil layer to surface soil (Jobbagy and Jackson 2004). More specific mechanisms may also be involved in the higher P uptake of *Solidago gigantea*, including mycorrhizal processes as documented for the closely related *S. canadensis* (Jin et al. 2004). *Solidago* might exploit its mycorrhizal symbiosis more effectively than the native grassland species. Zabinski et al. (2002) demonstrated the importance of mycorrhiza in affecting the competitive ability of an invasive forb over native grassland species and suggested that part of its ability to invade grasslands depends on the ability of its mycorrhizal symbiont to increase tissue P concentration. However, in our study, plant P concentration was lower for *Solidago* rather than for native plants.

Geochemical and biochemical processes involved in soil P transformations

Acid extractable inorganic P pools, i.e. Ca-bound P, can be depleted by plant species (Hedley et al. 1983; Zoysa et al. 1997). However, in our study, this fraction did not appear to be significantly affected by the invasive species (Tables 4 and 5) suggesting that it was under geochemical rather than biological control. The presence of *S. gigantea* increased all topsoil labile P pools, i.e. resin Pi, Bicarb-Pi and -Po, probably because sparingly soluble forms of soil P were mobilised by roots through increased efflux of H⁺ or exudation of organic acids (Hinsinger 2001 and references therein). The monospecific cover of *S. gigantea* significantly affected pH (ANOVA, Table 5) at one site, which probably indirectly affected the solubility

of P-containing complexes in solution (Lindsay 1979; Hinsinger, 2001). In addition to solubilization effect and because of its high affinity for di- and tri-valent cations, citrate and other organic acids can also displace P from humic-metal complexes, making it available for plant uptake (e.g. Frossard et al 1995; Jones 1998). However, these processes can only partly explain the increase of labile inorganic P as desorption and release of P by organic acids generally requires high concentrations of organic acids (Jones 1998).

Biologically mediated processes involved in soil P transformations

Differences in soil P status between control and invaded plots may also result from changes of soil microbiological activity as soil CO₂ release and phosphomonoesterase activities were significantly affected by the invasive species. Biological processes influence P distribution because most of the plant-available P is derived from soil organic matter (e.g. Cross and Schlesinger 1995; Frossard et al 1995, 2000). In our study, increased organic P content in topsoil may be related to higher litter inputs in invaded stands. An increase in bicarbonate extractable Po (Bicarb-Po), i.e. highly hydrolysable organic P is indicative of root exudation or high microbial activity and/or microbe and root turnover. The organic fraction extracted with hot concentrated (hcHCl-Po) was increased in topsoil under *Solidago* and may contain Po linked to particulate organic matter (POM). This macroorganic matter has been considered as an active organic matter pool that participates in the release of nutrients. Recent studies have indicated that the dynamics of POM can significantly influence P cycling in crop rotations with either legumes or natural fallow (Maroko et al. 1999; Salas et al. 2003), suggesting that P contained in this OM pool can have a significant influence on P availability. The linkage of labile Pi and organic P is reflected in enzymatic activities as organic P may contribute to plant P supply after being mineralized by enzymes. The influence of plant species on organic P mineralisation has been reported previously in various ecosystems (e.g.

Tarafdar and Claassen 1988; Giardina et al. 1995; Scott and Condron 2003). Labile inorganic P, i.e. resin- and bicarb-Pi was highly correlated to OH-Po and acid phosphomonoesterase activity.

Phosphomonoesterase activities in this study were in the range found in most other studies across a variety of ecosystems, 28-1670 $\mu\text{g NP g}^{-1} \text{ soil h}^{-1}$ (i.e. 0.2-12 $\mu\text{mol NP g}^{-1} \text{ soil h}^{-1}$; Nannipieri et al. 1978; Harrison 1983; Rojo et al. 1990; Bonmati et al. 1991; Amador et al. 1997). Phosphomonoesterase activities are correlated with OH-Po as expected because NaOH probably extracts compounds related to ortho-P monoesters (Robinson et al. 1998). Beside microorganisms, plant roots may represent an important source of acid phosphomonoesterase in soils, while they show no alkaline phosphatase activity (Juma and Tabatabai 1978). Alkaline phosphomonoesterase has been attributed to soil bacteria and fungi (Tarafdar and Claassen 1988) and soil microorganisms have been shown to be a major factor in controlling organic and inorganic P solution concentrations in temperate grassland topsoils (Seeling and Zasoski 1993). The plant species composition through root activity (exudation) is likely to determine critically the biomass, the diversity and the functional capacities of microbial species (Kourtev et al. 2002; Johnson et al. 2003; Chen et al. 2004). In addition to root exudates (Hinsinger 2001), differences in amount and quality (mineral nutrient concentration, C/N ratio, lignin and polyphenol contents) of organic debris between invasive species and resident vegetation may affect soil microbial community structure and function (Ehrenfeld 2003, Kourtev et al. 2002). Thus, in our studied situations, phosphomonoesterase activities were affected by the presence of the invasive species suggesting that the increase in the soil labile Pi pool under the cover of *Solidago gigantea*, was linked to enhanced Po mineralization. The significantly greater soil CO₂ release (indicative of soil respiration rate) and phosphomonoesterase activities may be due to enhanced soil microbial activity and diversity after invasion. Such a linkage between plant cover, soil microbiological activity and

soil P availability has been suggested by others (e.g. Tarafdar and Claassen 1988; Spears et al. 2001; Scott and Condron 2003).

Conclusions

It has been recognized that exotic invasive species can modify C and N cycling (Ehrenfeld 2003). Our results provide one of the first demonstrations of the influence of an invasive species on P availability and dynamics in soil. The results point out enhanced P dynamics in soil invaded by *Solidago gigantea*, probably through biologically mediated processes. However, this effect may vary with soil type and invading plant. The results also highlight that the impacts of exotic plants may not be apparent where only the total soil pool is considered, as observed by Ehrenfeld (2003) for C and N in her recent review. It is clear that a comprehensive understanding of the impacts of exotic species on soil requires a functional approach of the bioavailability of specific nutrients, including P.

Acknowledgements This study was supported by the Ministère de la Région de Bruxelles-Capitale (action “Research in Brussels”; scientific responsible: Claude Cocriamont-May) and was conducted in collaboration with the project INPLANBEL (Invasive Plants in Belgium: Patterns, Processes and Monitoring) funded by the Belgian Science Policy. The authors thank Evi Rossi and Ivan Nijs (University of Antwerp) for C-N analyses and Wolf Gruber (ULB) for analytical help.

References

- Amador JA, Glucksman AM, Lyons JB, Gorres JH (1997) Spatial distribution of soil phosphatase activity within a riparian forest. *Soil Sci.* 162:808-825.
- Bonmati M, Ceccanti B, Nannipieri P (1991) Spatial variability of phosphatase, urease, protease, organic carbon and total nitrogen in soil. *Soil Biol. Biochem.* 23:391-396.

- Bowman RA, Cole CV (1978) An exploratory method for fractionation of organic phosphorus from grassland soils. *Soil Sci.* 125:95–101.
- Chen CR, Condon LM, Davis MR, Sherlock RR (2004) Effects of plants species on microbial phosphorus and phosphatase activity in a range of grassland soils. *Biol. Fertil. Soils* 40:313-322.
- Cross AF, Schlesinger WH (1995) A literature review and evaluation of the Hedley fractionation: Applications to the biogeochemical cycle of soil phosphorus in natural ecosystems. *Geoderma* 64:197-214.
- Dalal RC (1977) Soil organic phosphorus. *Adv. Agron.* 29:83-117.
- Eivazi F, Tabatabai JM (1977) Phosphatases in soils. *Soil Biol. Biochem.* 9:167-172.
- Ehrenfeld JG (2003) Effect of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* 6:503-523.
- FAO/UNESCO (1990) Soil map of the world, Revised legend. World Soil Resources Report 60. FAO, Rome.
- Frossard E, Brossard M, Hedley M, Metherell A (1995) Reactions controlling the cycling of P in soils. In: Tiessen H (Ed) *Scope 54 - Phosphorus in the Global Environment - Transfers, Cycle's and Management.* John Wiley and Sons Ltd, Chichester, pp 107-138.
- Frossard E, Condon LM, Oberson A, Sinaj S, Fardeau JC (2000) Processes governing phosphorus availability in temperate soils. *J. Environ. Qual.* 29:15-23.
- Giardina CP, Huffman S, Binkley D, Caldwell BA (1995) Alders increase soil phosphorus availability in a Douglas-fir plantation. *Can. J. For. Res.* 25:1652-1657.
- Halstead RL, McKercher RB (1975) Biochemistry and cycling of phosphorus. In: Paul EA and McLaren AD (Eds) *Soil Biochemistry*, vol 3. Marcel Dekker, New York, pp 31-63.
- Harrison FA (1983) Relationship between intensity of phosphatase activity and physiochemical properties in woodland soils. *Soil Biol. Biochem.* 15:93-99.
- Hedley MJ and Stewart JWB (1982) Method to measure microbial phosphate in soils. *Soil Biol. Biochem.* 14:377–385.
- Hedley MJ, Nye PH, White RE (1983) Plant-induced changes in the rhizosphere of rape (*Brassica napus* Var. Emerald) seedlings. IV. The effect of rhizosphere phosphorus status on the pH, phosphatase activity and depletion of soil phosphorus fractions in the rhizosphere and on the cation-anion balance in the plants. *New Phytol.* 95:69-82.
- Hinsinger P (2001) Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. *Plant Soil* 237:173-195.
- Hobbie S (1992) Effects of plant species on nutrient cycling. *Trends Ecol. Evol.* 7:336-339.

- Hooper DU, Vitousek PM (1998) Effects of plant composition and diversity on nutrient cycling. *Ecol. Monogr.* 68:121-149.
- Jakobs G, Weber E, Edwards PJ (2004) Introduced plants of the invasive *Solidago gigantea* (Asteraceae) are larger and grow denser than conspecifics in the native range. *Divers. Distrib.* 10:11-19.
- Jin L, Gu Y, Xiao M, Chen J, Li B (2004) The history of *Solidago canadensis* invasion and the development of its mycorrhizal associations in newly-reclaimed land. *Func. Plant Biol.* 3:979-986.
- Jobbagy EG, Jackson RB (2004) The uplift of soil nutrients by plants: Biogeochemical consequences across scales. *Ecology* 85:2380-2389.
- John MK (1970) Colorimetric determination of phosphorus in soil and plant materials with ascorbic acid. *Soil Sci.* 109:214-220.
- Johnson D, Booth RE, Whiteley AS, Bailey MJ, Read DJ, Grime JP, Leake JR (2003) Plant community composition affects the biomass, activity and diversity of microorganisms in limestone grassland soil. *Eur. J. Soil Sci.* 54:671-677.
- Jones DL (1998) Organic acids in the rhizosphere - a critical review. *Plant Soil* 205:25-44.
- Juma NG, Tabatabai MA (1978) Distribution of phosphomonoesterases in soils. *Soil Sci.* 126:101-108.
- Kourtev PS, Ehrenfeld JG, Häggblom M (2002) Exotic plant species alter the microbial community structure and function in the soil. *Ecology* 83:3152-3166.
- Levine JM, Vilà M, D'Antonio C, Dukes JS, Grigulis K, Lavorel S (2003) Mechanisms underlying the impacts of exotic plant invasions. *P. Roy. Soc. Lond. B Bio.* 270:775-781.
- Lindsay WL (1979) *Chemical Equilibria in Soils*. John Wiley and Sons Ltd, New-York.
- López-Hernández D, Siegert G, Rodríguez JV (1986) Competitive adsorption of phosphate with malate and oxalate by tropical soils. *Soil Sci. Soc. Am. J.* 50:1460-1462.
- Maroko JB, Buresh RJ, Smithson PC (1999) Soil phosphorus fractions in unfertilized fallow-maize systems on two tropical soils. *Soil Sci. Soc. Am. J.* 63:320-326.
- Meyer AH, Schmid B (1999) Experimental demography of rhizome populations of establishing clones of *Solidago altissima*. *J. Ecol.* 87:42-54.
- Mitchell RJ, Marrs RH, Le Duc MG, Auld MHD (1997) A study of succession on lowland heaths in Dorset, southern England; changes in vegetation and soil chemical properties. *J. Appl. Ecol.* 34:1426-1444.
- Nannipieri P, Johnson RL, Paul EA (1978) Criteria for measurement of microbial growth and activity in soil. *Soil Biol. Biochem.* 10:223-229.

- Plichta W, Kuczyńska I, Rutkowski L (1997) The effect of American cherry (*Prunus serotina* Ehrh.) plantations on organic and humus horizons of cambic arenosols in a pine forest. *Ekol. Pol.- Pol. J. Ecol.* 45:385-394.
- Robinson JS, Johnston CT, Reddy KR (1998) Combined chemical and ³¹P-NMR spectroscopic analysis of phosphorus in wetland organic soils. *Soil Sci.* 163:705-712.
- Rojo MJ, Carcedo SG, Mateos MP (1990) Distribution and characterization of phosphatase and organic phosphorus in soil fractions. *Soil Biol. Biochem.* 22:169-174.
- Scott JT, Condon LM (2003) Dynamics and availability of Phosphorus in the rhizosphere of a temperate silvopastoral system. *Biol. Fertil. Soils* 39 (2):65-73.
- Salas AM, Elliott ET, Westfall DG, Cole CV, Six J (2003) The role of particulate organic matter in phosphorus cycling. *Soil Sci. Soc. Am. J.* 67:181-189.
- Seeling B, Zasoski RJ (1993) Microbial effects in maintaining organic and inorganic solution phosphorus concentrations in a grassland topsoil. *Plant Soil* 148:277-284.
- Spears JDH, Lajtha K, Caldwell BA, Pennington SB, Vanderbilt K (2001) Species effect of *Ceanothus velutinus* versus *Pseudotsuga menziesii*, Douglas-fir, on soil phosphorus and nitrogen properties in the Oregon cascades. *For. Ecol. Manag.* 149:205-216.
- StatSoft, Inc. (2003) STATISTICA (data analysis software system), version 6.1. www.statsoft.com.
- Tabatabai MA (1982) Soil Enzymes. In: Page AL, Miller RH and Keeney DR (Eds) *Methods of Soil Analysis, Part 2- Chemical and Microbiological Properties*. ASA-SSSA, Madison, WI, pp 903-915.
- Tarafdar JC, Claassen N (1988) Organic phosphorus compounds as a phosphorus source for higher plants through the activity of phosphatases produced by plant roots and microorganisms. *Biol. Fertil. Soils* 5:308-312.
- Tiessen H, Moir JO (1993) Characterization of available P by sequential extraction. In: Carter MR (Ed), *Soil Sampling and Methods of Analysis*. Lewis Publishers, Boca Raton, FL, pp 83-99.
- USDA-NRCS (2004) The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- Van Breemen N, Finzi AC (1998) Plant-soil interaction: ecological aspects and evolutionary implications. *Biogeochemistry* 42:1-19.
- Vitousek PM (1990). Biological invasion and ecosystem processes: towards an interaction of population biology and ecosystem studies. *Oikos* 57:7-13.

- Vitousek PM, D'Antonio CM, Loope LL, Rejmanek M, Westbrooks R (1997) Introduced species: a significant component of human-caused global change. *New Zeal. J. Ecol.* 21:1-16.
- Wagenitz G (1964) *Solidago* L. In: Hegi G. (Ed) *Illustrierte Flora von Mitteleuropa*, vol. VI. Carl Hanser, München, pp 16-29.
- Westover KM, Kennedy AC, Kelley SE (1997) Patterns of rhizosphere microbial community structure associated with co-occurring plant species. *J. Ecol.* 85:863-873.
- Zabinski CA, Quinn L, Callaway RM (2002) Phosphorus uptake, not carbon transfer, explains arbuscular mycorrhizal enhancement of *Centaurea maculosa* in the presence of native grassland species. *Funct. Ecol.* 16:758-765.
- Zoysa KN, Loganathan P, Hedley MJ (1997) A technique for studying rhizosphere processes in tree crops: soil phosphorus depletion around camellia (*Camellia japonica* L.) roots. *Plant Soil* 190:253-265.

Table 1 Description of the selected sites invaded by *Solidago gigantea*.

Site	Place in Belgium	Site type	Dominating plant species in uninvasion zones ^a
Sg1	Woluwe-St-Lambert	Waste land in urban area	<i>Dactylis glomerata</i> (G), <i>Arrhenatherum elatius</i> (G), <i>Heracleum sphondylium</i> (F), <i>Festuca rubra</i> (G), <i>Pastinaca sativa</i> (F), <i>Carex hirta</i> (G), <i>Agrostis stolonifera</i> (G), <i>Rubus sp.</i> (S)
Sg2	Kraainem	Rough grassland in urban area	<i>Agrostis stolonifera</i> (G), <i>Pulicaria dysenterica</i> (F), <i>Holcus lanatus</i> (G), <i>Epilobium hirsutum</i> (F), <i>Cirsium arvense</i> (F), <i>Leucanthemum vulgare</i> (F), <i>Festuca rubra</i> (G), <i>Achillea millefolium</i> (F)
Sg3	Saint-Ghislain	Rough grassland in nature conservation area	<i>Calamagrostis epigeios</i> (G), <i>Achillea millefolium</i> (F), <i>Epilobium angustifolium</i> (F), <i>Phragmites australis</i> (G), <i>Rubus sp.</i> (S), <i>Potentilla reptans</i> (F), <i>Cirsium arvense</i> (F)

^a Growth habit (USDA-NRCS 2004) in parentheses: G = graminoid; F = forb/herb; S = subshrub

Table 2 Properties of topsoil (0-10 cm) in invaded and uninvaded situations at three sites (means with standard deviations in parentheses, N = 6). *t*-tests: *ns*, not significant; *, *P* < 0.05.

Site		pH (H ₂ O)	CEC	CO ₂ release	Total C	Total N	CaCO ₃	Clay
			cmol kg ⁻¹	mg CO ₂ -C g ⁻¹ 14 days ⁻¹	%			
Sg1	Uninvaded	7.2 (0.2)	43.5 (18.8)	2.4 (0.3)	3.01 (0.8)	0.17 (0.05)	0.4 (0.1)	12.8 (3.4)
	Invaded	7.1 (0.1)	44.6 (10.7)	2.8 (0.3)	3.91 (0.9)	0.22 (0.04)	0.4 (0.2)	12.6 (2.1)
	<i>P</i> (<i>t</i> -test)	<i>ns</i>	<i>ns</i>	*	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
Sg2	Uninvaded	6.5 (0.4)	8.3 (0.9)	2.0 (0.2)	1.48 (0.16)	0.14 (0.01)	0.0	2.5 (0.5)
	Invaded	5.9 (0.3)	8.4 (1.2)	2.5 (0.4)	1.80 (0.54)	0.15 (0.02)	0.0	2.4 (0.4)
	<i>P</i> (<i>t</i> -test)	*	<i>ns</i>	*	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
Sg3	Uninvaded	7.1 (0.1)	39.2 (14.0)	2.7 (0.2)	9.36 (1.30)	0.36 (0.05)	1.4 (0.7)	17.0 (7.8)
	Invaded	7.2 (0.0)	59.9 (19.2)	2.9 (0.2)	8.51 (1.21)	0.36 (0.05)	1.3 (0.4)	14.1 (5.0)
	<i>P</i> (<i>t</i> -test)	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>

Table 3 Plant above-ground biomass and P concentration and uptake (Phytomass x P concentration) in invaded and uninvaded situations at the three sites (means with standard deviations in parentheses, N = 6). *t*-tests: *ns*, not significant; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Site		Phytomass	P concentration	P uptake
		g DM m ⁻²	mg g ⁻¹ DM	mg m ⁻²
Sg1	Uninvaded	357 (107)	2.4 (0.4)	848 (291)
	Invaded	917 (152)	1.9 (0.3)	1713 (227)
	<i>P</i> (<i>t</i> -test)	***	*	***
Sg2	Uninvaded	252 (126)	2.3 (0.2)	574 (287)
	Invaded	891 (102)	1.4 (0.2)	1211 (244)
	<i>P</i> (<i>t</i> -test)	***	***	**
Sg3	Uninvaded	438 (171)	2.0 (0.3)	878 (432)
	Invaded	760 (170)	2.0 (0.2)	1544 (461)
	<i>P</i> (<i>t</i> -test)	**	<i>ns</i>	*

Table 4 Comparison of P fractions (mg P kg⁻¹ soil) in invaded and uninvaded stands.

Site		Resin-Pi	Bicarb-		OH-		Acid-Pi	hc HCl-		Residual-P	Pi _(sum)	Po _(sum)	Pt _(sum)
			Pi	Po	Pi	Po		Pi	Po				
Sg1	Invaded	25.6	21.7	5.5	29.3	104.9	417.2	65.1	26.9	51.8	610.6	137.4	747.9
	Uninvaded	16.1	22.7	3.7	36.3	77.0	443.5	65.0	18.4	74.2	657.8	99.1	756.9
	<i>P</i> level ^a	ns ^b	ns	ns	ns	ns	ns	ns	ns	*	ns	ns	ns
Sg2	Invaded	28.1	23.0	12.5	42.0	44.6	55.1	53.1	20.6	59.7	261.0	77.7	334.1
	Uninvaded	19.3	19.0	11.0	38.1	36.8	57.0	51.8	15.8	60.1	245.3	63.6	308.9
	<i>P</i> level ^a	*	ns	ns	ns	ns	ns	ns	ns	ns	ns	*	ns
Sg3	Invaded	91.5	81.0	14.3	118.9	160.5	421.8	100.1	9.6	69.8	883.0	184.4	1066.9
	Uninvaded	61.2	57.9	11.3	98.0	121.0	469.7	96.6	6.1	98.0	881.	138.4	1019.8
	<i>P</i> level ^a	*	*	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns

^a Statistical significance for *t*-tests: ns, not significant; *, *P* < 0.05

^b not significant at 0.05 probability level, but note that here *P* = 0.06

Table 5 Results of two-way analyses of variance (ANOVA; df, *F*-ratio and probability levels) for effects of site, plant cover (uninvaded vs. invaded) and interactions on selected P fractions, phosphomonoesterase activities (Ac-PME and Alk-PME) and soil pH. Probability levels: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Source	df	Resin	Bicarb		OH		Acid	hcHCl		PME		pH
		Pi	Pi	Po	Pi	Po	Pi	Pi	Po	Ac	Alk	
Site (A)	2	76.7 ***	85.2***	26.2***	58.9***	11.5 ***	174.5***	28.5***	23.6***	21.0***	148.1***	52.8***
Plant (B)	1	15.4 ***	6.2*	4.8*	0.9NS	2.0 NS	1.6NS	0.1NS	9.9**	16.3***	30.9***	4.7*
A x B	2	2.9 NS	4.4*	0.2NS	1.6NS	0.3 NS	0.5NS	0.03NS	0.7NS	5.2*	10.0***	4.2*

Table 6 Pearson's correlation coefficients between selected P fractions and soil parameters (all sites pooled). Probability levels: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; NS, not significant. Coefficients higher than 0.70 are in bold.

	Bicarb-		OH-		PME		Soil pH	Total N	Total C
	Pi	Po	Pi	Po	Ac	Alk			
Resin-P	0.97***	0.51**	0.93***	0.71***	0.56***	0.52**	NS	0.72***	0.81***
Bicarb-Pi		0.44*	0.96***	0.73***	0.54***	0.53**	0.42*	0.78***	0.85***
Bicarb-Po			0.51**	NS	NS	NS	NS	NS	NS
OH-Pi				0.72***	0.50**	0.43*	NS	0.78***	0.82***
OH-Po					0.74***	0.70***	0.44*	0.60***	0.62***
Ac-PME						0.83***	0.45**	0.56***	0.58***
Alk-PME							0.76***	0.61***	0.64***
Soil pH								0.54***	0.58***
Total N									0.96***

Fig. 1 Acid (A) and alkaline (B) phosphomonoesterase (PME) activities in topsoil of invaded and uninvaded situations. Bars indicate standard deviation.

