Panarchy of an indigenous agroecosystem in the globalized market: The quinoa production in the Bolivian Altiplano

Thierry Winkel, Pierre Bommel, Marco Chevarría-Lazo, Geneviève Cortes, Carmen Del Castillo, Pierre Gasselin, François Léger, Juan-Peter Nina-Laura, Serge Rambal, Muriel 46 Tichit, et al.

► To cite this version:

Thierry Winkel, Pierre Bommel, Marco Chevarría-Lazo, Geneviève Cortes, Carmen Del Castillo, et al.. Panarchy of an indigenous agroecosystem in the globalized market: The quinoa production in the Bolivian Altiplano. Global Environmental Change, 2016, 39, pp.195-204. 10.1016/j.gloenvcha.2016.05.007 . ird-02381086

HAL Id: ird-02381086

<https://hal.ird.fr/ird-02381086>

Submitted on 27 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Panarchy of an indigenous agroecosystem in the globalized market: the quinoa production in the Bolivian Altiplano

Thierry Winkel^{a,*}, Pierre Bommel^b, Marco Chevarría-Lazo^c, Geneviève Cortes^d, Carmen Del Castillo^e, Pierre Gasselin^f, François Léger^g, Juan-Peter Nina-Laura^h, Serge Rambal^{i,j}, Muriel Tichit^g, Jean-François Tourrand^b, Jean-Joinville Vacher^k, Anaïs Vassas-Toral^d, Manuela Vieira-Pak^b, Richard Joffreⁱ

^aIRD, UMR 5175 CEFÉ, CNRS UM UPVM3 EPHE IRD, Montpellier, France; ^bCIRAD, UPR GREEN, Montpellier, France; ^cINRA, UMR MONA, Ivry-sur-Seine, France; ^dUniversité Paul Valéry, UMR 5281 ART-DEV, Montpellier, France; ^eFacultad de Agronomía, Universidad Mayor de San Andrés, La Paz, Bolivia; ^fINRA, UMR 951 Innovation, Montpellier, France; ^gUMR SADAPT, INRA, AgroParisTech, Université Paris-Saclay, Paris, France; ^hAgrónomos y Veterinarios sin Fronteras (AVSF), La Paz, Bolivia; ⁱCNRS, UMR 5175 CEFÉ, CNRS UM UPVM3 EPHE IRD, Montpellier, France; ^jDepartamento de Biología, Universidade Federal de Lavras, Lavras, MG, Brazil; ^kIRD, UMR 208 PALOC, IRD MNHN, Paris, France

* Corresponding author. Tel +33 467 613 328

Citation

Winkel T, Bommel P, Chevarría-Lazo M, Cortes G, Del Castillo C, Gasselin P, Léger F, Nina-Laura JP, Rambal S, Tichit M, Tourrand JF, Vacher JJ, Vassas-Toral A, Vieira-Pak M, Joffre R. 2016. Panarchy of an indigenous agroecosystem in the globalized market: the quinoa production in the Bolivian Altiplano. *Global Environmental Change* 39: 195–204. URL: <https://doi.org/10.1016/j.gloenvcha.2016.05.007>

Acknowledgements

This work was supported by the ANR (French National Research Agency) (ANR-06-PADD-011-EQUECO). M. Vieira-Pak benefited from an international grant of the McKnight Foundation (Programa Colaborativo de Investigación de Cultivos, CCRP-Andes-2011). We express our sincere gratitude to all the people, community authorities, agencies, and students who contributed to this project in Bolivia. Special thanks are due to the Programa de Investigación Estratégica en Bolivia (PIEB) for its valuable collaboration. Thanks are extended to the platform "Systèmes d'Information en Écologie" (CEFÉ) for assistance in remote sensing data analyses, and to Yvette Perfecto and Doyle McKey for helpful comments on early drafts of this work.

Abstract

Agricultural globalization is blamed for destructive impacts on small farms in developing countries. Yet, many local societies are proactive in the face of these changes and show high adaptive capacity. Investigating their transformations with an integrative perspective and enough hindsight may reveal some of the bases of their resilience and adaptive capacity. Using field data and the panarchy concept of resilience theory, we analyzed the territorial and social dynamics of quinoa growers' communities in southern Bolivia over the last four decades, a case study of regime shift in a poverty-stricken rural society which deliberately entered the global food market. Linking the dynamics of the household economy to the territorial and social subsystems over several decades, we gained insights into the interactions that shaped the rise of quinoa production in the region. We found that a vivid tradition of mobility allowing for pluriactivity on- and off-farm, combined with community self-governance, explains how local populations succeeded in articulating individual agency with collective control over their commons of land, seed resources, and social rules. Our vulnerability analysis points to landscape homogenization, social inequity, and increased dependence on external factors as potential sources of unsustainability. We conclude that, to cope with the changes of unprecedented magnitude they are facing, local producers should retain social cohesion and autonomous governance, without giving up on their heritage of mobility and economic redundancy. As regards theory, we identified cross-scale subsystem configurations critical for regime shifts, and confirm the value of panarchy in capturing complex socioecological dynamics.

Keywords

adaptive cycle – integrated assessment – land use change – rural livelihood strategies – transformative change

Highlights

- Rural societies in developing countries are not passive in the face of agricultural globalization.
- Hindsight and interdisciplinarity enlighten the bases of their proactivity and sustainability.
- We reconstructed the socioecological panarchy of Bolivian quinoa growers since 1970s.
- Commons governance and off-farm pluriactivity are critical to control latent unsustainability.
- Particular panarchy configurations give early-warning indicators for critical regime shifts.

1. Introduction

Quinoa is part of the diet of Andean populations for about 7000 years and has recently become a dietetic, organic, and gourmet commodity for Northern consumers (Bazile et al., 2015; Hellin and Higman, 2003). Family farmers of the southern Bolivian Altiplano initiated this change 40 y ago, but are now worried about the sustainability of their production. Soil erosion, biodiversity loss, and dietary degradation of local populations are pointed by development agents, researchers, and journalists, sometime using simplistic arguments and promoting questionable solutions (Winkel et al., 2012). A simple media campaign alerting consumers to emerging risks—be they real or not—could cause an immediate loss of income for local growers, and social and natural capital could be lost for rural Altiplano communities. Still, quinoa growers are quite efficient farmers and traders: producing a grain considered of little economic value until recently, they have succeeded in selling it, firstly to the neighboring market of Peru, and then to the international market in high-quality specialty foods (Giuliani et al., 2012; Laguna, 2011; Rojas et al., 2004). Their increasing efficiency was achieved by innovations such as partial crop mechanization or the incorporation in certified food-chains, combined with their traditional risk-coping strategy of mobility and off-farm pluriactivity (Kerssen, 2015; Padulosi et al., 2014; Walsh-Dilley, 2013). In fact, although quinoa growers produce for highly specialized export markets, they are not specialist farmers and are not even permanent dwellers in production areas. This is only one of the paradoxes that characterize quinoa production in this region. An agriculture booming under an extremely harsh environment, an organic and fair-trade production potentially threatening the local socioecosystem, and an opportunistic market searching for regulation: as will be detailed below, quinoa production is in many ways paradoxical.

Paradoxes reveal the complexity of reality, and sustainability science could benefit from analyzing and conceptualizing such situations to foster debates on its bases, methods, and goals, including political action and ethical controversies (Clark, 2007; Clark and Dickson, 2003; Müller, 2003; O'Connor, 2006; Vucetich and Nelson, 2010). Complexity is inherent to any human-environment interaction that includes various domains (ecological, social, economic, and cultural), scales (temporal and spatial), and goals (growth, equilibrium, or reduction) (Kajikawa, 2008). The concepts of adaptive cycles and panarchy help summarize part of this complexity. Adaptive cycles model living systems dynamics into successive phases of growth (or exploitation), conservation (or maturity), release (or collapse), and reorganization (or recycling) (Holling, 1973, 2001). Going a step further, panarchy models identify cross-scale interactions within nested hierarchies of adaptive cycles (Gunderson and Holling, 2002; Walker and Salt, 2006). These conceptual frameworks are powerful for articulating the complex structure of socioecosystems to their dynamics, transformability, and sustainability. They are relevant for many agricultural systems which show resilience in spite of harsh environment, weak rural policies, and low control of local stakeholders over foreign markets (Darnhofer et al., 2010; Darnhofer et al., 2016; Fraser and Stringer, 2009; Petrosillo et al., 2010; Salvia and Quaranta, 2015; Slaymaker, 2007).

This paper takes quinoa production in the Bolivian Altiplano as a test case to determine if adaptive cycle and panarchy models have empirical relevance for real sustainability issues. More specifically we seek to: (i) examine whether sustainability frameworks can yield insights into the paradoxes of an

ancestral food production recently entered into the globalized food market; (ii) identify emergent lessons for local management and policy in the world's major area of quinoa exportation; and (iii) identify scientific issues relevant for research on panarchy and sustainability.

2. Study area and theoretical backgrounds

2.1. Paradoxes, context, and drivers of quinoa success

Although quinoa is a specialized and highly valued export commodity, it is produced by part-time farmers frequently absent from the production areas. The rise in quinoa production triggered many emigrant people to return seasonally to their natal rural community for plowing, sowing and harvest. Here lies a major specificity of the social system sustaining quinoa production: local populations have high mobility and perform multiple farm- and non-farm activities in their originating rural communities and regional urban centers, respectively (Vassas-Toral, 2015c).

The natural environment in this region supports a flourishing export agriculture despite poor soils, cold desert climate, and high altitude. Crop lands, located 3,650–4,200 meters above sea level, receive an annual rainfall of 150–300 mm (regional south-north gradient), with frost on more than 200 days per year (Geerts et al., 2006; Pouteau et al., 2011). This harsh environment makes quinoa nearly the sole option for cropping. Regardless of its drought tolerance (Winkel et al., 2002), quinoa is unable to complete its growing cycle in that region with only the mean annual precipitation. Therefore, quinoa is grown in a 2-yr cycle (Joffre and Acho, 2008). After harvest, the land is left unseeded for a year but plowed so that the rainfall partly accumulates in the soil and becomes a water reserve available for the following year's crop (Giuliani et al., 2012)(**Supplementary Material A**). To our knowledge, quinoa is the only cash crop produced under low-input rainfed agriculture in extremely high, cold, and arid mountains.

Another paradox of quinoa is its status as a health food produced in small farms mostly under organic and fair-trade certification, but that might generate ecological and social ruptures. This would be opposite to the benefits expected from low-input organic agriculture in rural societies cultivating communal land supposedly governed by ancestral rules and traditional ecological knowledge. Researchers and journalists have blamed local growers for short-sightedness and individualism, but often with ill-founded arguments (Jacobsen, 2011; Winkel et al., 2014; Winkel et al., 2012). In fact, Bolivian farmers and decision makers are well aware of the rising vulnerability of their agroecosystem (Keressen, 2015). Local and individual initiatives at the origin of the expansion of the crop now seek collective regulations involving local authorities, growers' unions, rural development agencies, the central government, and international food industries. Moreover, and contrary to the common situation of rural poor excluded from their own natural capital (De Schutter, 2011; Kajikawa, 2008), Bolivian quinoa growers fully control access to their land resources and own on-farm selected seeds (Baudoin-Farah, 2009), and take opportunistic advantage of global trends in food demand.

2.2. Adaptive cycles and panarchy

Adaptive cycles model the dynamics of a living system into successive phases of growth or exploitation (r), conservation (K), release (Ω), and reorganization (α), which provides for system

renewal (Holling, 1973, 2001) (**Fig. 1A**). Cycle speed and amplitude are controlled by two system properties: connectedness (or degree of organization) and potential (or amount of resources). Connectedness and potential are defined depending on the ecological, economic, or sociocultural specificities of the considered adaptive cycle. Novelty in adaptive cycles may appear by the end of the reorganization (α) phase, when low internal control (low connectedness) allows for external opportunities (or "chance events") to nucleate and open unexpected options for a new growth phase. Adaptive cycles should not be seen as 4-phase sequences "repeating themselves over history" (Bunce et al., 2009): depending on the adaptive capacity of the system, transitions may occur between virtually all phases (Walker and Salt, 2006). To avoid the deterministic connotation of the term "cycle", we adopt hereafter the expression "adaptive loop" used by various authors (e.g. Delcourt and Delcourt, 2004; Marsh, 2016).

A complete panarchy model emerges when a nested hierarchy of adaptive loops is identified and cross-scale interactions are recognized (Gunderson and Holling, 2002; Walker and Salt, 2006) (**Fig. 1B**). A special case of cross-scale interaction, called "revolt" connection, occurs when a small-scale subsystem enters the Ω phase while the surrounding system is in late K phase (intermediate loop in **Fig. 1B**). This uniform and rigid intermediate-scale system is vulnerable to the propagation of a subsystem collapse ("revolt" arrow from the lower-scale subsystem at Ω) which may trigger a major catastrophic event. A typical example of such dynamics is the propagation of a local, small-scale fire (an Ω release phase) within a monospecific and even-aged forest in an advanced K phase. The renewal of the intermediate-scale subsystem will depend on the resources stored at the upper-scale subsystem ("remember" arrow). Typically, the recovery of a burned area will draw upon the seed bank and the surviving species present in the surrounding territory.

3. Methods

To gain insights into the social and ecological drivers of the agricultural sustainability in the world's leading area of quinoa production, a research-action project was implemented from 2007 to 2010 in rural communities of the departments of Potosi and Oruro in the southern Altiplano of Bolivia. This participative multidisciplinary research involved specialists in agroecology, agronomy, geography, and socioeconomics, with quinoa grower families and local stakeholders (workers in development agencies, quinoa growers' unions, local authorities, etc.). The results of this sustainability assessment are detailed in Vassas-Toral (2015c), Vieira-Pak (2015) and Winkel et al. (2015), and only relevant information will be retained here.

3.1. Socioeconomic surveys

The patterns of mobility and pluriactivity among quinoa growers from the study area were characterized in a socioeconomic survey conducted in 2007 and 2008 on 149 households from five rural communities (San Juan, Chilalo, Otuyo, Candelaria, Palaya). Using interviews and livelihood stories, we retraced the residential and professional trajectories of 170 members of quinoa grower families (**Supplementary Material B**). Different types of spatial mobilities were considered: (i) the migration of individuals, defined as a temporary or permanent change of residence, including mobility of bi-residential migrants, (ii) the seasonal and circular mobility of migrants linked to the agricultural

activity in their community. Collected data were structured by mapping the successive residencies of each surveyed individual, and retracing the chronology of his/her life sequences (see Vassas-Toral, 2015abc, for detailed results). A second survey using semi-directive interviews was conducted in 2007 on 36 households from six communities (Buena Vista, Chacoma, Chilalo, Jirira, Playa Verde Murmuntani, San Agustin) focusing on the household economy and the farming practices of quinoa growers (see Acosta-Alba, (2007) for detailed results).

3.2. Land use mapping

Cropland expansion in the last five decades was mapped on a sample of six communities representative of quinoa production in the study area, with altitudes ranging from 3700 to 4100 masl, at distances from the shores of the Salar of Uyuni varying from 2 to 40 km (**Supplementary Material C**). Some localities are mostly flat plains (Otuyo, Candelaria), while others are mostly sloping (Chilalo) or with mixed plain/slope landscapes (Capura, Chacoma, Palaya).

Cropland mapping was based on aerial photographs and high-resolution satellite images. Early panchromatic aerial photographs (Instituto Geográfico Militar, Bolivia) taken from 1961 to 1963 show the croplands before the quinoa crop expansion. EROS satellite images (United States Geological Survey, USA) dated from 27/04/1972 coincide with the initial phase of quinoa expansion in the Salar region. SPOT satellite images (Centre National d'Études Spatiales, France) taken between 2005 and 2008 show the phase of fast expansion of the quinoa crop. After digitalization, the ground resolution of the aerial photographs was 2 m. Satellite images had a ground resolution of 10 m for EROS data, and 20 to 28.5 m for SPOT (depending of the year of acquisition). Image distortions due to the sensors and the uneven ground surface were rectified using a digital elevation model (SRTM 90m). All the images were georeferenced in the UTM 19 S projection, with the datum Provisional South American 1956 mean. Croplands were identified visually in the images as polygons of bare soil areas, clearly differentiated from the steppic vegetation of the surrounding pastures and non-agricultural land. After extraction with the ENVI 4.2 software, the cropland polygons identified were superimposed to a digital elevation model with a 20-m resolution using the ArcGIS/ArcView software. This allowed examining the altitudinal distribution of crop areas in each community, as well as their temporal changes across successive images.

3.3. Vulnerability assessment

Sustainability science needs vulnerability assessments integrating the human and environmental components of socioecosystems. The "vulnerability scoping diagram" presented by Polsky et al. (2007) provides multidisciplinary research teams with a common framework for organizing dissimilar information about the three dimensions of vulnerability in human-environment systems exposed to hazards, namely: exposure, sensitivity and adaptive capacity (Polsky et al., 2007; Turner II et al., 2003). We used this conceptual diagram as a guide for the collection of data, the definition of indicators and the specification of a set of hazards and outcomes relevant for quinoa production in the study area (**Supplementary Material D**).

4. Results and discussion

4.1 Vulnerability assessment of southern Altiplano agriculture

Our multidisciplinary research in the Bolivian Altiplano identified quinoa field expansion as the major indicator of environmental change in the study area (Winkel et al., 2015). Quantitative mapping of six communities shows that total cropland area from 1963 to 2006 increased by 332%, mostly in flat plains suitable for mechanization (**Fig. 2**). National statistics show that the total quinoa crop area in Bolivia jumped from approximately 22,600 ha in 1963 to 42,431 ha in 2006, and 173,960 ha in 2014 (FAOSTAT, 2016). Quinoa yield variations and inequity in household incomes provide further vulnerability indicators. A socioeconomic survey of 36 households in 2007 revealed two consequences of the recent land use changes: (i) mean areas ($\pm$ CV%) of individual quinoa fields increased from 3.1 ha $\pm$ 4% on slopes, to 9.3 ha $\pm$ 39% in plains; and (ii) mean quinoa yield ($\pm$ CV) decreased from 0.91 t·ha⁻¹ $\pm$ 13% on manually cultivated slopes to 0.60 t·ha⁻¹ $\pm$ 13% on mechanically cultivated plains. The increased CV in field area reflects a growing inequity among growers. Wealthy growers and tractor owners capture and cultivate much larger land areas than poor growers. This growing inequity affects household monetary incomes, which ranged from 200–18,000 US\$ per year among the 36 surveyed families. Independent studies report similar socioeconomic changes (Astudillo and Aroni, 2012; Félix and Villca, 2009; Kerksen, 2015; Ormachea and Ramirez, 2013). The grain yield decrease reflects the sensitivity of quinoa production to mechanized cropping in plains. Soil overexploitation has been claimed as the cause of reduced yield (Félix and Villca, 2009; Jacobsen, 2011). However, other causes could include poor seed germination in mechanically sowed fields, increased frost risks in lowlands, and pest proliferation in large monocultures. These and other ecological and social indicators of the local agroecosystem show its vulnerability to recent land use and crop management changes. Still, quinoa growers adaptively respond to new market opportunities, and display commitment to diversified livelihood strategies, despite the unprecedented success of the global quinoa market (Kerksen, 2015; Vassas-Toral, 2015c; Walsh-Dilley, 2013).

Grasping the ins and outs of the ongoing mutation of the quinoa socioecosystem requires adding to this static review a dynamic scaling of the structures and processes driving the system. To this end, recent changes in social context, household economy, and territorial patterns, were plotted in adaptive loop models, and we analyzed where the growth, conservation, release and reorganization phases of the adaptive loops fit these changes.

4.2. Adaptive loops in the southern Bolivian Altiplano

4.2.1. Social subsystem

Historically, human demography in this highland region has been characterized by pendular migrations to nearby Andean valleys or to the Pacific coast to exchange salt and agricultural goods or to temporarily escape severe drought or frost in the Altiplano (Murra, 1984). Cycles of prosperity and collapse in Bolivian and Chilean mining industries since the 19th century also contributed to population movements in the region (Platt, 1995). From 1950 onward, distant cities attracted ever more people in search of job opportunities and new lifestyles (Vassas-Toral, 2015c; Walsh-Dilley, 2013). Yet, the most recent Bolivian census (2012) shows an incipient repopulation dynamics, suggesting a possible re-peasantization of the southern Altiplano (Kerksen, 2015). At the same time, conflicts concerning

land access are proliferating in the communities because migrants with double residency in the city and the community claim land access rights inherited from their parents (**Supplementary Material B**). The national census registers these migrants in the community, which allows them to reaffirm local land rights. These migrations are generally seasonal or temporary, and quantifying whether rural communities are effectively regaining population or undergoing a demographic decline remains a complex issue (Laguna, 2011; Vassas-Toral, 2015c).

Regulations intended to guide local land access and use are also complex and ambiguous. The 1952 agrarian reform had little impact in this inhospitable region neglected by great landlords (Laguna, 2011; Walsh-Dillely, 2013), and a multiplicity of rules have subsequently piled up. Local rules and customs transmitted and controlled by traditional indigenous authorities co-exist with national laws enacted by the central government. As regards crop production, collective norms resulting from local consensus compete with food industry controls, foreign trade regulations, and intellectual property rights on quinoa varieties. Most of these rules are not uniformly applied in the region due to low rates of acceptance and lack of practical implementation. Some development agencies designate this plethora of non-uniform regulations as a "legal vacuum" (Félix and Villca, 2009). Thus, plotting population demography and social regulations as the respective potential and connectedness axes of the social adaptive loop in the region (**Fig. 3A**), the current situation appears to be a long-lasting reorganization (α) phase, with emigrants temporarily mixing with returning people, and with mixed rules of land management still not implemented. Notably, reorganization does not occur here after a collapse (Ω) phase of ancient local societies, but rather as progressive adjustments of their population dynamics and land management rules.

4.2.2. Household subsystem

Amid this varying social framework, the quinoa economy has flourished rapidly in the region. Quinoa as a cash crop expanded during 1970–1980 from the southern border to the western and northern borders of the Salar of Uyuni. This expansion was initiated in response to: i) massive job cuts in the mining and public sectors due to structural adjustment plans, ii) increasing demand for quinoa in neighboring Peru, a country much more populated than Bolivia and where urban consumers appreciated quinoa, which was not the case in Bolivia at that time. In 1970, a Belgian development aid program donated tractors to local, poor communities, a case of "chance event" which favored that initial phase of commercial production for the national and Peruvian markets (Kerssen, 2015; Laguna, 2011). In the mid 1980's, Bolivian quinoa growers opportunistically responded to new, increased demand from North America and Europe for gluten-free, protein-rich, organic food, often under fair-trade label. These new markets did not supplant the Peruvian market, which accounted for up to 40% of total Bolivian quinoa exports by the 2000's, though mostly as unofficial trade (Gandarillas et al., 2015; Laguna, 2002; Laguna, 2011; Rojas, 2011). Thus, local quinoa growers beneficially serve a diversity of quinoa market niches. Yet, this success did not spur them to fully specialize in quinoa production. The vast majority maintain off-farm activities and temporary migration (Supplementary Material B: Table B-1 and reported individual cases), with a few even keeping on with llama and sheep rearing, despite lower economic profitability. Diversified household trajectories result from this redundancy in more or less rewarding activities involving different family members (Laguna, 2002;

Vassas-Toral, 2015c). This risk-coping strategy may prove to be useful under the market distortions observed since 2014, when Peru began to supplant Bolivia in the international quinoa market (Mercaderó, 2015). If the household economic loop is plotted with quinoa income and market connection as the potential and connectedness axes respectively (**Fig. 3B**), then quinoa growers of the southern Bolivian Altiplano appear to follow multiple trajectories between the growth (r) and conservation (K) phases, resolutely engaged in commercial production but avoiding connecting their household activities exclusively to that opportunistic venture. Partial back loops exist, and moving from one trajectory to another is relatively easy (Vassas-Toral, 2015c; Vieira-Pak, 2015). For example, if a quinoa grower opted for the specialized certified market, he can easily move to the less constraining and almost equally rewarding conventional market, or may choose to invest more effort in off-farm activities (Ofstehage, 2012). However, achieving this flexibility is not easy for the poorest households (single mothers or elderly single people) due to the lack of human and financial capital. The current variety of household trajectories also reflects past social and economic disparities, which are exacerbated by quinoa commercialization (Vassas-Toral, 2015c; Vieira-Pak, 2015).

4.2.3. Territorial subsystem

This subsystem appears more constrained than the household subsystem. A single quinoa harvest is the product of a 2-year cycle and, thus, represents a doubled land area: the current cultivated field, plus the plowed fallow waiting for the next year's crop (**Supplementary Material A**). Quinoa expansion pushed herbaceous and shrub species to marginal lands. These wild plants hardly recolonize the fallow due to their slow dynamics and the depletion of soil seed banks (Joffre and Acho, 2008). Conversion of pastures into quinoa fields thus results in an essentially irreversible change in land use, with a striking loss of landscape patchiness. Bare fallows are exposed to strong and frequent windstorms. The newly converted quinoa fields are mostly located in flat lowlands prone to frost due to cold air drainage at night (Pouteau et al., 2011). Frost events in 2007 and 2008 revealed the vulnerability of lowland production with 2008 referred to as a tipping point in the quinoa market (Núñez de Arco, 2015). Production was low due to these frost events but commercial demand remained high, which caused the quinoa price to increase by 175% from 2006 to 2008 (Gabriel, 2013). These high prices encouraged farmers to continue taking the risk of cultivating quinoa in the lowlands.

Most of the areas apt for mechanized cropping have already been converted into quinoa fields, and many communities can no longer respond to the continuing demand for new croplands. A decline of soil fertility in the region has been denounced (Félix and Villca, 2009; Jacobsen, 2011), allegedly showing that the ecosystem's carrying capacity has been exceeded. However, this opinion has not been corroborated by any *in situ* study, and the evidence used to support it consists of gross regional or national grain yield statistics, which mix concomitant effects of climate, soil, pests, and poor seed germination in mechanized crops (Walsh-Dilley, 2013; Winkel et al., 2012). While soil degradation remains debatable, land conversion to quinoa cropping thus appears as the most reliable indicator of the ecosystem's carrying capacity reaching an upper limit. Land conversion resulted from the decision of local producers to favour quinoa at the expense of the economically less-rewarding components of their agricultural system, namely pastures and potato crops (Laguna, 2011; Walsh-Dilley, 2013). After a sustained growth phase (r), the territorial adaptive loop appears locked into an ending conservation

phase (*K*) with contiguous quinoa monocultures resulting in minimum landscape patchiness (maximum field connectedness, x-axis), and the maximum available land area (natural capital, y-axis) already converted to cropland in most communities (**Fig. 3C**).

4.3. *Panarchy of the quinoa's rise*

4.3.1. *Nested adaptive loops*

A hierarchy linking the adaptive loops for social, household, and territorial subsystems is organized with respect to time and population scales (**Fig. 4**). The household loop occupies the lower scales since it primarily refers to individual decisions regarding annual on-farm and off-farm activities. Typically, household dynamics occur on a 1- to 6-month scale, e.g. to choose temporary migration or which parcel to plow. The territorial loop is located at the larger landscape scale, with a minimum 2-year cycle for field crop, and a grossly decadal dynamics for land use changes and natural vegetation regeneration. Next, the social loop operates from the community to the nation and from decadal to multidecadal periods corresponding to background demographic trends and land access/land use regulations. Demographic and institutional changes commonly show slower dynamics than do ecological changes (Young, 2010). In fact, the social loop could be extended up to the global level, as the current international market directly affect the local agrarian dynamics through the commercial demand and trade regulations (Gabriel, 2013; Giuliani et al., 2012; Rojas, 2011).

The phases currently reached by each adaptive loop show that the territorial component is locked in a potentially vulnerable conservation phase (*K*) between household and social loops (**Fig. 4B**). Yet, a "revolt" connection (see **Fig. 1B**) amplified by this broad-scale vulnerability seems unlikely, as the lower-scale household economic loop is far from the release phase (Ω). The diversified household livelihood strategies, associated to rural-urban mobility (**Supplementary Material B**) (Laguna, 2011; Vassas-Toral, 2015c) prevent any locking into a specialized commodity chain. If commodity chains were to experience sudden collapse, it would trigger the abandonment of croplands—a release phase Ω in the territorial adaptive loop—but without catastrophic impact on households which opportunistically maintain their economies between growth (*r*) and conservation (*K*) extremes by repeated migration and shifting between farm- and non-farm activities (**Fig. 3B**).

Because the trajectories of many agroecosystems lack of certain phases or appear stuck in particular stages of the adaptive loop, some authors call into question the usefulness of Holling's model in resilience studies (Bunce et al., 2009). Also, it has been argued that Gunderson and Holling's theory is more adequate to describe ecological processes than to capture adaptive responses related to human decisions (Fraser and Stringer, 2009). Both critiques should be relativized since: i) adaptive loops are not deterministic 4-phase sequences: the panarchy theory states that alternative paths and shortcuts might occur within adaptive loops, some of which open the way towards resilience (Gunderson and Holling, 2002; Walker and Salt, 2006); besides, stagnation in one adaptive loop of the system does not impede rapid changes in other components, as found comparing the social subsystem to the other two components in our case study; ii) in agroecosystems in particular, alternative paths of resilience depend strongly on human decisions: our analysis of quinoa growers' strategies demonstrates that these decisions are adequately captured by an adaptive loop model.

Thus, contrary to the suggestion of Fraser and Stringer (2009), it seems not justified to restrain the panarchy approach to the environmental component of the socio-ecosystem. In fact, the main interest of the panarchy theory lies in its capacity to encompass environmental as well as social, economic, and institutional dynamics: disaggregating the environmental and human components of change may be necessary in an early phase of analytical assessment but then, ordering them across nested hierarchies to reveal their interdependencies is what makes the value of this transdisciplinary and relational approach (Darnhofer et al., 2016; Kajikawa, 2008; Scholz and Steiner, 2015; Sundstrom et al., 2014).

4.3.2. Retroactive search for causalities

A retroactive examination of quinoa panarchy can provide insight into recent events (Westley, 2002) and identify circumstances, actors, and organizations that shaped the historical development of quinoa production in southern Bolivia. At the end of the 1960s, crop production and pastoralism provided low income due to the lack of markets, conducting a growing part of the local population to frequent and prolonged migration to cities and other countries in search of non-farm activities. Thus, the territorial and household subsystems were in a reorganization (α) phase but with still uncertain prospects as regards local development (**Fig. 4A**). The declining rural population was ageing, which caused local institutions and social rules to lose vitality. The social subsystem found itself caught between collapse (Ω) and reorganization (α) phases (**Fig. 4A**). The wealthiest among the permanent rural dwellers were those who owned a substantial number of sheep and llamas and had greatest access to communal pastures. Rural migrants with savings from off-farm activities were also in a favorable position to assert land usufruct rights which families always retain in their natal communities. Both categories of households had the potential to invest capital in new economic activities. As regards the territorial land resources at that time, there was little competition for land access and little control on land use due to the low interest in farming activities. In 1970, these circumstances enabled some individuals to respond opportunistically to newly available tractors and the continuous Peruvian demand for quinoa. New quinoa fields were opened in common pastures in flat lowlands, easily mechanizable, lacking individual property rights, and considered of little value. The social, territorial, and household subsystems were then positioned in similar reorganization (α) phases, ready to produce a cascade of new patterns across the hierarchy of adaptive loops—viz the panarchy—of the local quinoa production (**Fig. 4A**, dashed line). In the following years, however, the dynamics of the three adaptive loops rapidly diverged. This led to the present situation (**Fig. 4B**) with multiple household activities and adjustments (loop E between r and K), maximum pressure on land resources (loop T near K), and prolonged experimentation and uncertainty in social and institutional issues (loop S near α).

4.4. Lessons for local development

The rise of quinoa production in southern Bolivia began 40 years ago. Large changes have occurred in society, agrotechnology, and economics which preclude a mere return to traditional agricultural models. The transition from ancestral manual cropping to mechanized cropping exemplifies an agricultural revolution (Mazoyer and Roudart, 2006) that triggered an unprecedented landscape

change. As in other rural societies in the Andes (Lennox and Gowdy, 2014; Zimmerer, 2013), new challenges regarding land management have emerged, which require new visions for a sustainable future.

Panarchy highlights the importance of individual initiatives for adaptability of quinoa growers. Mobility and pluriactivity are key components of the household's functional redundancy, a concept that date back to an ancestral economic mobility (Núñez Atencio and Hall, 1982; Saignes, 1995), and has been adjusted to current technical and trading innovations. With a singularity however: for the first time in centuries, local agriculture is an economically attractive enterprise, which stimulates seasonal returns of migrants to their natal communities (Walsh-Dille, 2013). Since growing quinoa entails a hereditary access to communal land, it enables autonomy to local populations, which is radically different from selling their labor force in cities or mining industries. This change was quickly assimilated by rural dwellers who then adopted a patrimonial logic and tried to hoard the largest possible cropland area, within the bounds of the customary land tenure rules. This quantitative patrimonialization rapidly reached the limits of available space, and a qualitative patrimonialization is now needed to improve cropping practices, and negotiate comprehensive agreements for using common land resources.

Thus far, the social system also contributed to changes in quinoa production, though in an ambiguous way. Heirs of a tradition of self-governance and rotating public responsibilities, local authorities and quinoa growers' unions did not exert any strong social control on their own members until recently. This behavior enabled the free transformation of an agroecosystem with manual cropping and long fallows to mechanized monocultures and short fallows. The prevalence of oral rules during the early transformation may have promoted these changes. Ambiguity in the social control system also appears to be part of a logic of the local institutions to avoid contradictory commitments to the traditional subsistence system and the market-oriented system. Rivalry between local authorities and quinoa growers' unions could emerge, the latter relaying the top-down market demand for international norms of sustainable production, and the former engaging in a bottom-up renovation of local ancestral rules governing their commons. Both adaptability and transformability are essential for resilience (Walker and Salt, 2006). In the southern Bolivian Altiplano, the household adaptability and the social transformability may result in contradicting influences (see subsequent discussion on adaptation traps). These properties rely primarily on the capacity of the institutions controlling common resources to evolve by self-transformation, which requires social cooperation and commitment to avoid centralized regulation or privatization (Vollan and Ostrom, 2010).

The observed territorial vulnerability suggests a central role for the landscape structure in determining the agroclimatic risks faced by quinoa growers. Frost risks persist in the region despite climate warming trends (Pouteau et al., 2011; Rambal et al., 2015), and monocultures in vast lowland areas may worsen drought, pest outbreaks, and soil degradation. Local agronomic institutes are conducting participatory research on these issues to design innovative land management practices (Bonifacio et al., 2014). Numerous communities have begun using integrated organic pest management and planting living fences against wind erosion (Lino et al., 2014). These agroecological practices promoted by farmers' unions and NGOs illustrate social learning and innovation drawn from

local experimentations (Knight and Meffe, 1997). Pilot experiments on new payment for ecosystem
service schemes are another example of collective action to conserve local quinoa biodiversity
(Narloch et al., 2015).

The southern Bolivian Altiplano socioecosystem conforms to the conditions for adaptive land
management—persistency, change and uncertainty (Holling, 1973). Persistency of the local society
relies on mobility and flexibility of the growers' families, self-governance of their communities and
organizations, and material and symbolic territorial valorization. Changes in land use and household
activities enable land transformation as new opportunities appear. This illustrates how local
populations cope with, rather than confront, uncertainty due to fluctuations in climate or economy
(Walsh-Dille, 2013). Similar synergies between off-farm migration, governance of the commons and
innovation leading to social and ecological resilience have been described in smallholders
communities engaged in agricultural intensification in other regions of Bolivia (Zimmerer, 2013). In
other Andean regions, however, the sustainability of smallholder agriculture may rely on different
bases, depending on local settings of land tenure, social organization, opportunities for off-farm
activities and market access (Lennox and Gowdy, 2014; Padulosi et al., 2014; Sietz et al., 2012).

Although quinoa growers tried to adaptively manage their territorial resources, there are several
indicators of latent unsustainability. In some communities, vast quinoa monocultures have generated
conflicts over land access (Laguna, 2011; Vassas-Toral, 2015c). This partly resulted from familial and
social agreements that permit families to cultivate without being present in the community. These are
the same agreements that support household adaptability and territorial transformability, illustrating an
adaptation trap in which incremental adjustments and economic redundancies delay local stakeholder
awareness of agroecosystem vulnerabilities (van Apeldoorn et al., 2011). All stakeholders do not
share the same capacity to access social and territorial resources, or to activate multiple options of
economic redundancies. The poorest families with the lowest social capital also are the most
disadvantaged in competition for new croplands. In this context, livelihood diversification further
increases social inequity among families. An undesirable system could thus emerge in which
household adaptability generates a trap that could threaten social equity and ecological sustainability.

To face such complex issues, development NGOs, state or private services offer their expertise.
Indeed, the rationality of local populations is no longer the only one that applies in the Bolivian
altiplano: new foreign actors introduce divergent opinions and practices, which increases the quinoa
socioecosystem complexity. Now, complexity also can cause unsustainability (Fisk and Kerherve,
2006). Some of the new foreign actors promote short-sighted interests and irreversible technological
innovations like irrigation or certified seeds, with a lack of integrative vision which has great potential
to worsen the situation. These actors reinforce the command-and-control approach to natural resource
management (Briggs, 2003; Holling and Meffe, 1996). If this approach is followed, local societies
would become increasingly dependent on risky technologies, bank credits, and rationales external to
the local social and environmental context. Southern Altiplano populations are fluent with foreign
markets and different cultures, as part of their inherited lifestyle based on active mobility and trade
exchange (Stern, 1995). Yet, the unprecedented magnitude of these emerging socioeconomic
interactions raises concerns (Lennox and Gowdy, 2014). Local policy-makers and stakeholders should

cautiously evaluate these new forms of interference and dependence before making decisions that bind the region and its inhabitants to a future they may not desire. Such a loss of autonomy would challenge the goals of good living and food sovereignty recently integrated into the Bolivian constitution (Cockburn, 2013; Kerksen, 2015; Mercado et al., 2016).

5. Conclusion

This study reminds us that panarchy is more than just the adaptive loop of a specific subsystem: a panarchy is essentially a *dynamic hierarchy of several subsystems*—environmental and human—that interact over nested scales of time and space. Using this multi-dimensional and dynamic perspective, we were able to elucidate part of the sustainability of quinoa production in the southern altiplano of Bolivia during the last four decades, firstly by identifying the characteristics of the households, the environment, and the communities at stake. Then, by examining interactions between these nested components, we found that the adhesion of local populations to self-governance and the tradition of temporary migration, instead of causing social stagnation, had indeed stimulated the rapid transformation of their territory. This is explained by processes operating at different scales: a limited social control in the initial phase of crop expansion, the interdiction for foreigners to access land resources in the communities, and household's pluriactivity as a continued risk-coping strategy despite the boon of the quinoa trade.

While nested hierarchies of scales may be relatively trivial to detect (household, community, state, etc.), their driving interactions are less obvious because the underlying dynamics often require time to develop and become apparent. Conceptual efforts should be made to enrich the typology of these interactions and discover new cross-scale connections beyond the well-described configurations of revolt or memory (Walker and Salt, 2006) (**Fig. 1B**). The synchronous alignment of nested subsystems in reorganization phase (α) found in this study is an example of such critical configurations (**Fig. 4A**). It reveals a situation open to innovations, chance events, and regime shifts cascading across subsystems. This alignment pattern within a complex socioecosystem may provide an early-warning indicator, a form of intrinsically cross-scale transition signal (Scheffer et al., 2009). Despite such conceptual advances, scholars and decision makers should accept that no panacea exists for modeling or governing socioecosystems (Anderies and Janssen, 2013; Brock and Carpenter, 2007; Ostrom, 2007). These are controlled simultaneously by fast and slow variables, and go through incessant co-adaptation processes that continuously change their structure, function, and interrelationships, with inevitable dysfunctions (Reynolds et al., 2007; Stafford Smith et al., 2007). In such systems, static equilibria never arise and top-down command-and-control governance fails (Holling and Meffe, 1996). Instead, continuous adaptive learning and polycentric governance copes with uncertainties and opportunities, particularly in agricultural or natural resource management (Anderies and Janssen, 2013; Darnhofer et al., 2010). Because panarchy models help identifying people and organizations responsible for the choices and consequences of development policies, an ethical issue rises up: topics of equity, sustainability, and autonomy of local societies stem from these choices (Vucetich and Nelson, 2010), and thus challenge the social utility of scholarly research (Thompson, 2008). Agroecosystems offer fertile ground for such ethical questioning in sustainability science.

References

- Acosta-Alba, I., 2007. Durabilité des systèmes de production de l'altiplano sud bolivien : quels équilibres entre élevage et agriculture ? AgroParisTech, Université Paris 7, Paris, France, 44 p.
- Alvarez-Flores, R., Winkel, T., Degueldre, D., Del Castillo, C., Joffre, R., 2014a. Plant growth dynamics and root morphology of little-known species of *Chenopodium* from contrasted Andean habitats. *Botany* 92, 101-108. <http://dx.doi.org/10.1139/cjb-2013-0224>
- Alvarez-Flores, R., Winkel, T., Nguyen-Thi-Truc, A., Joffre, R., 2014b. Root foraging capacity depends on root system architecture and ontogeny in seedlings of three Andean *Chenopodium* species. *Plant and Soil* 380, 415-428. <http://dx.doi.org/10.1007/s11104-014-2105-x>
- Anderies, J.M., Janssen, M.A., 2013. Robustness of social-ecological systems: implications for public policy. *Policy Studies Journal* 41, 513-536. <http://dx.doi.org/10.1111/psj.12027>
- Astudillo, D., Aroni, G., 2012. Livelihoods of quinoa producers in southern Bolivia, in: Giuliani, A., Hintermann, F., Rojas, W., Padulosi, S. (Eds.), *Biodiversity of Andean grains: balancing market potential and sustainable livelihoods*. Bioversity International, Rome, Italy, pp. 78-145. [http://www.bioversityinternational.org/index.php?id=19&user_bioversitypublications_pi1\[showUId\]=7303](http://www.bioversityinternational.org/index.php?id=19&user_bioversitypublications_pi1[showUId]=7303)
- Baudoin-Farah, A., 2009. Evaluación y perspectivas del mercado de semillas certificadas de quinua en la región del Salar de Uyuni en el Altiplano Sur de Bolivia. Master Thesis, AgroParisTech, Département SIAFEE, Paris, France, 35 p.
- Bazile, D., Bertero, D., Nieto, C., 2015. State of the art report on quinoa around the world in 2013. FAO (Santiago de Chile) y CIRAD, (Montpellier, Francia), Rome, Italy, 589 p. URL: <http://www.fao.org/583/a-i4042e.pdf>
- Bonifacio, A., Aroni, G., Villca, M., Ramos, P., Alcon, M., Gandarillas, A., 2014. The current and potential role of q'ila q'ila (*Lupinus* spp.) in quinoa sustainable production systems. *Revista de Agricultura (Bolivia)* 54, 11-18. http://www.proinpa.org/publico/REVISTA_54_INGLES.pdf
- Briggs, S., 2003. Command and control in natural resource management: revisiting Holling and Meffe. *Ecological Management & Restoration* 4, 161-162. <http://dx.doi.org/10.1046/j.1442-8903.2003.00151.x>
- Brock, W.A., Carpenter, S.R., 2007. Panaceas and diversification of environmental policy. *Proceedings of the National Academy of Sciences of the United States of America* 104, 15206-15211. <http://www.pnas.org/content/104/39/15206.abstract>
- Bunce, M., Mee, L., Rodwell, L.D., Gibb, R., 2009. Collapse and recovery in a remote small island: a tale of adaptive cycles or downward spirals? *Global Environmental Change* 19, 213-226. <http://dx.doi.org/10.1016/j.gloenvcha.2008.11.005>
- Clark, W.C., 2007. Sustainability science: a room of its own. *Proceedings of the National Academy of Sciences of the United States of America* 104, 1737-1738. <http://dx.doi.org/10.1073/pnas.0611291104>

Clark, W.C., Dickson, N.M., 2003. Sustainability science: the emerging research program.

622 Proceedings of the National Academy of Sciences of the United States of America 100, 8059-
8061. <http://dx.doi.org/10.1073/pnas.1231333100>

624 Cockburn, J., 2013. Bolivia's food sovereignty & agrobiodiversity: undermining the local to strengthen
the state?, International Conference "Food Sovereignty: A Critical Dialogue", September 14-15,
626 2013, Yale University, USA. URL: [https://www.tni.org/en/briefing/bolivias-food-sovereignty-](https://www.tni.org/en/briefing/bolivias-food-sovereignty-agrobiodiversity)
agrobiodiversity.

628 Darnhofer, I., Bellon, S., Dedieu, B., Milestad, R., 2010. Adaptiveness to enhance the sustainability of
farming systems. A review. Agronomy for Sustainable Development 30, 545-555.

630 <http://dx.doi.org/10.1051/agro/2009053>

Darnhofer, I., Lamine, C., Strauss, A., Navarrete, M., 2016. The resilience of family farms: towards a
632 relational approach. Journal of Rural Studies 44, 111-122.

<http://dx.doi.org/10.1016/j.jrurstud.2016.01.013>

634 De Schutter, O., 2011. The green rush: the global race for farmland and the rights of land users.

Harvard International Law Journal 52, 503-559. [http://www.harvardilj.org/2011/07/issue_52-2_de-](http://www.harvardilj.org/2011/07/issue_52-2_de-schutter/)
636 [schutter/](http://www.harvardilj.org/2011/07/issue_52-2_de-schutter/)

Delcourt, P.A., Delcourt, H.R., 2004. Prehistoric native Americans and ecological change. Human
638 ecosystems in Eastern North America since the pleistocene. Cambridge University Press,
Cambridge, UK.

640 FAOSTAT 2016. Quinoa production. Statistics Division, Food and Agriculture Organization of the
United Nations, Rome, Italy. Retrieved 31/03/2016: http://faostat3.fao.org/search/*/E

642 Félix, D., Villca, C., 2009. Quinoa y territorio: Experiencias de acompañamiento a la gestión del
territorio y a la autogestión comunal en la zona Intersalar del altiplano boliviano. VSF-CICDA-
644 Ruralter, Lyon, France. <https://www.avsf.org/public/posts/564/quinoa-y-territorio.pdf>

Fisk, D.J., Kerherve, J., 2006. Complexity as a cause of unsustainability. Ecological Complexity 3,
646 336-343. <http://dx.doi.org/10.1016/j.ecocom.2007.02.007>

Fraser, E.D.G., Stringer, L.C., 2009. Explaining agricultural collapse: macro-forces, micro-crises and
648 the emergence of land use vulnerability in southern Romania. Global Environmental Change 19,
45-53. <http://dx.doi.org/10.1016/j.gloenvcha.2008.11.001>

650 Gabriel, A., 2013. Royal Quinoa reigning the world: from food of the poor to world's superfood.
Bachelor Thesis, Van Hall Larenstein - Applied University of Science, Wageningen, The
652 Netherlands, 80 p. http://mercadero.nl/wp-content/uploads/Final_Thesis_AG_FTM_1.7.pdf

Gandarillas, A., Rojas, W., Bonifacio, A., Ojeda, N., 2015. Quinoa in Bolivia: the PROINPA
Foundation's perspective, in: Bazile, D., Bertero, D., Nieto, C. (Eds.), State of the art report on
quinoa around the world in 2013. FAO, Santiago, Chile, pp. 344-377. URL:
656 <http://www.fao.org/343/a-i4042e.pdf>

Geerts, S., Raes, D., Garcia, M., Del Castillo, C., Buytaert, W., 2006. Agro-climatic suitability mapping
658 for crop production in the Bolivian Altiplano: a case study for quinoa. Agricultural and Forest
Meteorology 139, 399-412. <http://dx.doi.org/10.1016/j.agrformet.2006.08.018>

Giuliani, A., Hintermann, F., Rojas, W., Padulosi, S., 2012. Biodiversity of Andean grains: balancing market potential and sustainable livelihoods. Bioversity International, Rome, Italy, 198 p.
[http://www.bioversityinternational.org/index.php?id=19&user_bioversitypublications_pi1\[showUId\]=7303](http://www.bioversityinternational.org/index.php?id=19&user_bioversitypublications_pi1[showUId]=7303)

Gunderson, L.H., Holling, C.S., 2002. Panarchy: understanding transformations in human and natural systems. Island Press, Washington (DC), USA, 507 p.

Hellin, J., Higman, S., 2003. Quinoa and food security, in: Hellin, J., Higman, S. (Eds.), Feeding the Market: South American Farmers, Trade and Globalization. Intermediate Technology Development Group (ITDG) Publishing, London, UK, pp. 131-168.

Holling, C.S., 1973. Resilience and stability of ecological systems. Annual Review of Ecology and Systematics 4, 1-23. <http://dx.doi.org/10.1146/annurev.es.04.110173.000245>

Holling, C.S., 2001. Understanding the complexity of economic, ecological, and social systems. Ecosystems 4, 390-405. <http://dx.doi.org/10.1007/s10021-001-0101-5>

Holling, C.S., Meffe, G.K., 1996. Command and control and the pathology of natural resource management. Conservation Biology 10, 328-337. <http://dx.doi.org/10.1046/j.1523-1739.1996.10020328.x>

Jacobsen, S.E., 2011. The situation for quinoa and its production in southern Bolivia: from economic success to environmental disaster. Journal of Agronomy and Crop Science 197, 390-399.
<http://dx.doi.org/10.1111/j.1439-037X.2011.00475.x>

Joffre, R., Acho, J., 2008. Quinua, descanso y tholares en el sur del Altiplano Boliviano. Hábitat 75, 38-41. <http://www.lidema.org.bo/>

Kajikawa, Y., 2008. Research core and framework of sustainability science. Sustainability Science 3, 215-239. <http://dx.doi.org/10.1007/s11625-008-0053-1>

Kerssen, T.M., 2015. Food sovereignty and the quinoa boom: challenges to sustainable re-peasantisation in the southern Altiplano of Bolivia. Third World Quarterly 36, 489-507.
<http://dx.doi.org/10.1080/01436597.2015.1002992>

Knight, R.L., Meffe, G.K., 1997. Ecosystem management: agency liberation from command and control. Wildlife Society Bulletin 25, 676-678. <http://research-legacy.arch.tamu.edu/epsru/pdf/plan641/Plan%20PDFs%202/Plan%20PDFs%202/Knight%20and%20Meffe.pdf>

Laguna, P., 2002. Competitividad, externalidades e internalidades, un reto para las organizaciones económicas campesinas: la inserción de la Asociación Nacional de Productores de Quinua en el mercado mundial de la quinua. Debate Agrario (Perú) 34, 95-169.
<http://www.cepes.org.pe/debate/debate34/04-articulo-da34.pdf>

Laguna, P., 2011. Mallas y flujos: acción colectiva, cambio social, quinua y desarrollo regional indígena en los Andes Bolivianos. School of Social Sciences, Wageningen University, Wageningen, The Netherlands, 516 p. <http://edepot.wur.nl/188049>

Lennox, E., Gowdy, J., 2014. Ecosystem governance in a highland village in Peru: facing the challenges of globalization and climate change. Ecosystem Services 10, 155-163.
<http://dx.doi.org/10.1016/j.ecoser.2014.08.007>

- Lino, V., Olivera, J., Saravia, R., Quispe, R., Gandarillas, E., Crespo, L., 2014. Mass dissemination of the Integrated Pest Management (IPM) strategy for quinoa. *Revista de Agricultura (Bolivia)* 54, 68-72. http://www.proinpa.org/publico/REVISTA_54_INGLES.pdf
- Marsh, E.J., 2016. The disappearing desert and the emergence of agropastoralism: an adaptive cycle of rapid change in the mid-Holocene Lake Titicaca Basin (Peru–Bolivia). *Quaternary International*, in press. <http://dx.doi.org/10.1016/j.quaint.2015.12.081>
- Mazoyer, M., Roudart, L., 2006. A history of world agriculture from the neolithic age to the current crisis. Translated by James H. Membréz. Earthscan Publ., London, UK.
- Mercadero, 2015. European quinoa market continues fast growth while prices have dropped. Mercadero, Wageningen, The Netherlands; Retrieved 2015/12/23: <http://mercadero.nl/category/quinoa-market-analysis/>
- Mercado, G., Hjortsø, C.N., Kledal, P.R., 2016. Public procurement for school breakfasts in the Bolivian Altiplan: governance structures enabling smallholder inclusion. *Journal of Rural Studies* 44, 63-76. <http://dx.doi.org/10.1016/j.jrurstud.2016.01.004>
- Müller, A., 2003. A flower in full blossom?: ecological economics at the crossroads between normal and post-normal science. *Ecological Economics* 45, 19-27. [http://dx.doi.org/10.1016/S0921-8009\(02\)00256-2](http://dx.doi.org/10.1016/S0921-8009(02)00256-2)
- Murra, J.V., 1984. Andean societies. *Annual Review of Anthropology* 13, 119-141. <http://dx.doi.org/10.1146/annurev.an.13.100184.001003>
- Narloch, U., Drucker, A.G., Pascual, U., 2015) What role for cooperation in conservation tenders? Paying farmer groups in the High Andes. *Land Use Policy*, in press. <http://dx.doi.org/10.1016/j.landusepol.2015.09.017>
- Núñez Atencio, L., Hall, H., 1982. Análisis de dieta y movilidad en un campamento arcaico del Norte de Chile. *Bulletin de l'IFEA* 11, 91-113. <http://www.ifeanet.org/publicaciones/boletines/11%283-4%29/91.pdf>
- Núñez de Arco, S., 2015. Quinoa's calling, in: Murphy, K.M., Mataguihan, J. (Eds.), *Quinoa: Improvement and Sustainable Production*. John Wiley & Sons, Hoboken, NJ, USA, pp. 211-226.
- O'Connor, M., 2006. The "Four Spheres" framework for sustainability. *Ecological Complexity* 3, 285-292. <http://dx.doi.org/10.1016/j.ecocom.2007.02.002>
- Ofstehage, A., 2012. The construction of an alternative quinoa economy: balancing solidarity, household needs, and profit in San Agustín, Bolivia. *Agriculture and Human Values* 29, 441-454. <http://dx.doi.org/10.1007/s10460-012-9371-0>
- Ormachea, E.S., Ramirez, N.F., 2013. Propiedad colectiva de la tierra y producción agrícola capitalista: el caso de la quinua en el Altiplano Sur de Bolivia. Centro de Estudios para el Desarrollo Laboral y Agrario - CEDLA, La Paz, Bolivia. <http://mercadero.nl/wp-content/uploads/libro-quinua-2013-pdf.pdf>
- Ostrom, E., 2007. A diagnostic approach for going beyond panaceas. *Proceedings of the National Academy of Sciences of the United States of America* 104, 15181-15187. <http://dx.doi.org/10.1073/pnas.0702288104>

- Padulosi, S., Amaya, K., Jäger, M., Gotor, E., Rojas, W., Valdivia, R., 2014. A holistic approach to enhance the use of neglected and underutilized species: the case of andean grains in Bolivia and Peru. *Sustainability* 6, 1283-1312. <http://www.mdpi.com/2071-1050/6/3/1283>
- Petrosillo, I., Zaccarelli, N., Zurlini, G., 2010. Multi-scale vulnerability of natural capital in a panarchy of social-ecological landscapes. *Ecological Complexity* 7, 359-367. <http://dx.doi.org/10.1016/j.ecocom.2010.01.001>
- Platt, T., 1995. Ethnic calendars and the market interventions among the Ayllus of Lipes during the nineteenth century, in: Larson, B., Harris, O., Tandeter, E. (Eds.), *Ethnicity, markets, and migration in the Andes: at the crossroads of history and anthropology*. Duke University Press, Durham, USA, pp. 259-296.
- Polsky, C., Neff, R., Yarnal, B., 2007. Building comparable global change vulnerability assessments: the vulnerability scoping diagram. *Global Environmental Change* 17, 472-485. <http://dx.doi.org/10.1016/j.gloenvcha.2007.01.005>
- Pouteau, R., Rambal, S., Ratte, J.-P., Gogé, F., Joffre, R., Winkel, T., 2011. Downscaling MODIS-derived maps using GIS and boosted regression trees: the case of frost occurrence over the arid Andean highlands of Bolivia. *Remote Sensing of Environment* 115, 117-129. <http://dx.doi.org/10.1016/j.rse.2010.08.011>
- Rambal, S., Ratte, J.P., Mouillot, F., Winkel, T., 2015. Trends in quinoa yield over the southern Bolivian altiplano: lessons from climate and land-use projections, in: Murphy, K.M., Matanguihan, G.J. (Eds.), *Quinoa Improvement and Sustainable Production*. John Wiley & Sons, Hoboken, NJ, USA, pp. 47-62.
- Reynolds, J.F., Stafford Smith, D.M., Lambin, E.F., Turner II, B.L., Mortimore, M., Batterbury, S.P.J., Downing, T.E., Dowlatabadi, H., Fernández, R.J., Herrick, J.E., Huber-Sannwald, E., Jiang, H., Leemans, R., Lynam, T., Maestre, F.T., Ayarza, M., Walker, B., 2007. Global desertification: building a science for dryland development. *Science* 316, 847-851. <http://dx.doi.org/10.1126/science.1131634>
- Rojas, W., 2011. Quinoa, an ancient crop to contribute to world food security. FAO, Oficina Regional para America Latina y el Caribe, Santiago, Chile, 58 p. http://www.fao.org/fileadmin/templates/aig2013/res/en/cultivo_quinua_en.pdf
- Rojas, W., Soto, J.L., Carrasco, E., 2004. Study on the social, environmental and economic impacts of quinoa promotion in Bolivia. PROINPA Foundation, La Paz, Bolivia, 84 p. http://www.underutilized-species.org/Documents/PUBLICATIONS/quinoa_case_study_es.pdf
- Saignes, T., 1995. Indian migration and social change in seventeenth-century Charcas, in: Larson, B., Harris, O., Tandeter, E. (Eds.), *Ethnicity, markets, and migration in the Andes: at the crossroads of history and anthropology*. Duke University Press, Durham, USA, pp. 167-195.
- Salvia, R., Quaranta, G., 2015. Adaptive cycle as a tool to select resilient patterns of rural development. *Sustainability* 7, 11114. <http://www.mdpi.com/2071-1050/7/8/11114>
- Scheffer, M., Bascompte, J., Brock, W.A., Brovkin, V., Carpenter, S.R., Dakos, V., Held, H., van Nes, E.H., Rietkerk, M., Sugihara, G., 2009. Early-warning signals for critical transitions. *Nature* 461, 53-59. <http://dx.doi.org/10.1038/nature08227>

- Scholz, R.W., Steiner, G., 2015. Transdisciplinarity at the crossroads. *Sustainability Science* 10, 521-526. <http://dx.doi.org/10.1007/s11625-015-0338-0>
- Sietz, D., Mamani Choque, S.E., Lüdeke, M.K.B., 2012. Typical patterns of smallholder vulnerability to weather extremes with regard to food security in the Peruvian Altiplano. *Regional Environmental Change* 12, 489-505. <http://dx.doi.org/10.1007/s10113-011-0246-5>
- Slaymaker, O., 2007. The potential contribution of geomorphology to tropical mountain development: the case of the MANRECUR project. *Geomorphology* 87, 90-100. <http://dx.doi.org/10.1016/j.geomorph.2006.06.044>
- Stafford Smith, D.M., McKeon, G.M., Watson, I.W., Henry, B.K., Stone, G.S., Hall, W.B., Howden, S.M., 2007. Learning from episodes of degradation and recovery in variable Australian rangelands. *Proceedings of the National Academy of Sciences of the United States of America* 104, 20690-20695. <http://www.pnas.org/cgi/content/abstract/104/52/20690>
- Stern, S.J., 1995. The variety and ambiguity of native Andean intervention in European colonial markets, in: Larson, B., Harris, O., Tandeter, E. (Eds.), *Ethnicity, markets, and migration in the Andes: at the crossroads of history and anthropology*. Duke University Press, Durham, USA, pp. 73-100.
- Sundstrom, S.M., Angeler, D.G., Garmestani, A.S., García, J.H., Allen, C.R., 2014. Transdisciplinary application of cross-scale resilience. *Sustainability* 6, 6925-6948. <http://dx.doi.org/10.3390/su6106925>
- Thompson, P.B., 2008. Agrarian philosophy and ecological ethics. *Science and Engineering Ethics* 14, 527-544. <http://dx.doi.org/10.1007/s11948-008-9094-1>
- Turner II, B.L., Kasperson, R.E., Matson, P.A., McCarthy, J.J., Corell, R.W., Christensen, L., Eckley, N., Kasperson, J.X., Luers, A., Martello, M.L., Polsky, C., Pulsipher, A., Schiller, A., 2003. A framework for vulnerability analysis in sustainability science. *Proceedings of the National Academy of Sciences of the United States of America* 100, 8074-8079. <http://dx.doi.org/10.1073/pnas.1231335100>
- van Apeldoorn, D.F., Kok, K., Sonneveld, M.P.W., Veldkamp, T.A., 2011. Panarchy rules: rethinking resilience of agroecosystems, evidence from dutch dairy-farming. *Ecology and Society* 16, 39. URL: <http://www.ecologyandsociety.org/vol16/iss31/art39/>
- Vassas-Toral, A., 2011. Ruralité et agriculture au prisme des mobilités spatiales. *L'Altiplano Sud de Bolivie à l'heure du boom de la quinoa*. Université Paul Valéry - Montpellier III, Montpellier, France, 345 p. + annexes. <https://tel.archives-ouvertes.fr/tel-00668162/>
- Vassas-Toral, A., 2015a. Movilidades de los productores de quinua y dinámicas territoriales en el Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), *Racionalidades campesinas en los Andes del Sur: reflexiones en torno al cultivo de la quinua y otros vegetales andinos*. EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 231-280.
- Vassas-Toral, A., 2015b. Movilidades, recursos de tierra y sistema de producción agrícola en el Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), *Racionalidades campesinas en los Andes del Sur: reflexiones en torno al cultivo de la quinua y otros vegetales andinos*. EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 281-321.

- Vassas-Toral, A., 2015c. *Partir et cultiver: essor de la quinoa, mobilités et recompositions rurales en Bolivie*. IRD Éditions, Marseille, France.
- Vieira-Pak, M., 2015. El auge de la quinoa: visiones de futuro en el Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), *Racionalidades campesinas en los Andes del Sur: reflexiones en torno al cultivo de la quinoa y otros vegetales andinos*. EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 323-353.
- Vollan, B., Ostrom, E., 2010. Cooperation and the commons. *Science* 330, 923-924.
<http://dx.doi.org/10.1126/science.1198349>
- Vucetich, J.A., Nelson, M.P., 2010. Sustainability: virtuous or vulgar? *BioScience* 60, 539-544.
<http://dx.doi.org/10.1525/bio.2010.60.7.9>
- Walker, B.H., Salt, D.A., 2006. *Resilience thinking: sustaining ecosystems and people in a changing world*. Island Press, Washington DC, USA.
- Walsh-Dille, M., 2013. Negotiating hybridity in highland Bolivia: indigenous moral economy and the expanding market for quinoa. *The Journal of Peasant Studies* 40, 659-682.
<http://dx.doi.org/10.1080/03066150.2013.825770>
- Westley, F., 2002. The devil in the dynamics: adaptive management on the front lines, in: Gunderson, L.H., Holling, C.S. (Eds.), *Panarchy: understanding transformations in human and natural systems*. Island Press, Washington DC, USA, pp. 333-360.
- Winkel, T., Alvarez-Flores, R., Bertero, D., Cruz, P., Del Castillo, C., Joffre, R., Peredo Parada, S., Sáez Tonacca, L., 2014. Calling for a reappraisal of the impact of the quinoa expansion on agricultural sustainability in the Andean highlands. *Idesia (Chile)* 32, 95-100.
<http://dx.doi.org/10.4067/S0718-34292014000400012>
- Winkel, T., Alvarez-Flores, R.A., Bommel, P., Bourliaud, J., Chevarria Lazo, M., Cortes, G., Cruz, P., Del Castillo, C., Gasselin, P., Joffre, R., Léger, F., Nina Laura, J.P., Rambal, S., Rivière, G., Tichit, M., Tourrand, J.F., Vassas-Toral, A., Vieira-Pak, M., 2015. The southern altiplano of Bolivia, in: Bazile, D., Bertero, D., Nieto, C. (Eds.), *State of the art report on quinoa around the world in 2013*. FAO/Cirad, Santiago, Chile, pp. 362-377. URL: <http://www.fao.org/363/a-i4042e.pdf>
- Winkel, T., Bertero, H.D., Bommel, P., Chevarria Lazo, M., Cortes, G., Gasselin, P., Geerts, S., Joffre, R., Léger, F., Martinez Avisa, B., Rambal, S., Rivière, G., Tichit, M., Tourrand, J.F., Vassas-Toral, A., Vacher, J.J., Vieira-Pak, M., 2012. The sustainability of quinoa production in southern Bolivia: from misrepresentations to questionable solutions. Comments on Jacobsen (2011, *J. Agron. Crop Sci.* 197: 390–399). *Journal of Agronomy and Crop Science* 198, 314-319.
<http://dx.doi.org/10.1111/j.1439-037X.2012.00506.x>
- Winkel, T., Méthy, M., Thénot, F., 2002. Radiation-use efficiency, chlorophyll fluorescence and reflectance indices, associated with ontogenic changes in water-limited *Chenopodium quinoa* leaves. *Photosynthetica* 40, 227-232. <http://dx.doi.org/10.1023/A:1021345724248>
- Young, O.R., 2010. Institutional dynamics: resilience, vulnerability and adaptation in environmental and resource regimes. *Global Environmental Change* 20, 378-385.
<http://dx.doi.org/10.1016/j.gloenvcha.2009.10.001>

Zimmerer, K.S., 2013. The compatibility of agricultural intensification in a global hotspot of smallholder
agrobiodiversity (Bolivia). Proceedings of the National Academy of Sciences of the United States
of America 110, 2769-2774. <http://dx.doi.org/10.1073/pnas.1216294110>

FIGURE CAPTIONS

Fig. 1. Stylized representations of an adaptive loop (A) and a panarchy (B) showing "revolt" and "remember" connections (adapted from *Panarchy*, Gunderson & Holling, eds. 2002)

Fig. 2. Cropland expansion in the southern Bolivian altiplano from 1963 to 2006. (A) Palaya community. (B) Altitudinal distribution of total cropland area in 6 communities of the southern Bolivian Altiplano.

Fig. 3. Adaptive loops for the social subsystem (A), household subsystem of quinoa growers (B), and territorial subsystem (C) in the southern Bolivian Altiplano.

Fig. 4. Quinoa panarchy in the southern Bolivian Altiplano from the 1970's (A) to the present (B). Dotted lines connect the relative positions of the adaptive loops of household economy (E), territorial land resources (T), and social regulations (S) in each time period. Black arrows in (B) indicate changing alternatives in household economy.

SUPPLEMENTARY MATERIAL CAPTIONS

Supplementary Material A. The quinoa crop cycle in the southern altiplano of Bolivia.

Supplementary Material B. Mobility and migration among quinoa growers.

Supplementary Material C. Localisation and cropland area in six communities of the southern altiplano of Bolivia from 1963 to the 2000's.

Supplementary Material D. The Vulnerability Scoping Diagram.

FIGURES

Fig. 1. Stylized representations of an adaptive loop (A) and a panarchy (B) showing "revolt" and "remember" connections (adapted from *Panarchy*, Gunderson & Holling, eds. 2002).

FIGURES

Fig. 2. Cropland expansion in the southern Bolivian altiplano from 1963 to 2006. (A) Palaya community. (B) Altitudinal distribution of total cropland area in 6 communities of the southern Bolivian Altiplano.

Fig. 3. Adaptive loops for the social subsystem (A), household subsystem of quinoa growers (B), and territorial subsystem (C) in the southern Bolivian Altiplano.

912

914 **Fig. 4.** Quinoa panarchy in the southern Bolivian Altiplano from the 1970's (A) to the present (B).
Dotted lines connect the relative positions of the adaptive loops of household economy (E), territorial
916 land resources (T), and social regulations (S) in each time period. Black arrows in (B) indicate
changing alternatives in household economy.

918

SUPPLEMENTARY MATERIAL

Supplementary Material A. The quinoa crop cycle in the southern altiplano of Bolivia

Cultivation in that region has to cope with two major climatic constraints: a large frost period from April to October and a short rainy season between December and March. In fact, frost events may occur even during the rainy season, and the mean annual rainfall (lower than 250 mm) does not cover the water requirements of a complete quinoa crop cycle. Thus, the quinoa crop cycle necessarily takes place over a two year period including a long period of plowed fallow with bare soil.

The crop calendar begins in Year 1 with the plowing of the land area planned to be harvested in Year 2. Plowing is done in January-February in order to facilitate rainfall infiltration and refilling of the soil water reserve. Nowadays, in the plains, this work is most often mechanized. During the austral winter (from May to August), soil evaporation is reduced. Soils are generally sandy, and soil moisture in the subsurface soil layer in late August is sufficient for early sowing and seed germination (R. Joffre, unpublished data). Early sowing is advantageous as it allows for harvesting the crop before the March-April frost period, but it is also risky as the probability of a severe frost event damaging the quinoa seedlings in September-October is non-negligible. Quinoa seeds placed on the humid soil layer at a 10-20 cm depth germinate and fastly develop deep rooting (Alvarez-Flores et al., 2014a; Alvarez-Flores et al., 2014b). The growing season lasts till March-April in Year 2 and the harvest has to be done before the main frost period (corresponding to the austral autumn-winter). During the post-harvest period, a large part of the soil surface remains without any vegetation cover.

	Year 1												Year 2											
Month	J	F	M	A	M	J	J	A	S	O	N	D	J	F	M	A	M	J	J	A	S	O	N	D
Climate																								
rainy season																								
main frost risks																								
Land use	plowed field								cultivated field						fallow									
Vegetation	bare soil										5 – 40%						< 10%							
Cropping																								
plowing																								
sowing																								
crop growth																								
harvest																								

Cited references

- Alvarez-Flores, R., Winkel, T., Degueldre, D., Del Castillo, C., Joffre, R. 2014a. Plant growth dynamics and root morphology of little-known species of *Chenopodium* from contrasted Andean habitats. Botany 92, 101-108. DOI: 10.1139/cjb-2013-0224.
- Alvarez-Flores, R., Winkel, T., Nguyen-Thi-Truc, A., Joffre, R. 2014b. Root foraging capacity depends on root system architecture and ontogeny in seedlings of three Andean *Chenopodium* species. Plant Soil 380, 415-428. DOI 410.1007/s1104-11014-12105-x.

Supplementary Material B. Mobility and migration among quinoa growers

This section shows patterns of mobility and pluriactivity among quinoa growers from the southern Altiplano of Bolivia. We considered several types of spatial mobilities: (i) the migration of individuals, defined as a temporary or permanent change of residence, including mobility of bi-residential migrants, (ii) the seasonal and circular mobility of migrants linked to the agricultural activity in their community.

A socioeconomic survey was conducted in two years (2007, 2008) on 149 households from 5 rural communities (San Juan, Chilalo, Otuyo, Candelaria, Palaya). Using interviews and livelihood stories, we reconstructed the residential and professional trajectories of 170 members of quinoa grower families (Vassas-Toral, 2011, 2015a, b, c). The results show (i) the extension of the migration area of 139 migrants people between 1934 and 2008 (Fig. B-1) and the diversity of the activities practised by male and female migrants in this period (Table B-1) , (ii) two cases illustrating the individual trajectories of migrant people from these communities (Figs. B-2, 3), (iii) two cases illustrating the circular mobility trajectory linked to the agricultural activity (Fig. B-4, 5). To preserve confidentiality, the names of the interviewed people have been changed.

Fig. B-1. Extension of the migration area of 139 migrant people from 5 rural communities of the southern altiplano of Bolivia between 1934 and 2008.

Source: Interviews and livelihood stories, 2007-2008. Elaboration: A. Vassas-Toral and S. Coursière.

The mapping of the migrations of the 170 surveyed individuals (139 of them effectively migrating) reveals more than 400 events between 1934 and 2008, with urban and rural destinations in Bolivia, Chile and Argentina (Fig. B-1). In 2008, the main destinations of migration were big cities of Bolivia (La Paz, Oruro, Cochabamba) and Chile (Iquique). **Table B-1** (below) shows the professional activities practised by 139 migrating people, differentiating male and female migrants (data are percent of interviewees who declared having practised the activity in at least one of their migrations):

Economic sector	Division	Status and activity	Males (%)	Females (%)
primary	agriculture	informal sector employee: farmhand, lumberer	31	6
		self-employed worker: farmer, tractor driver	10	14
	mining	informal sector employee: miner	44	0
secondary	building industry	informal sector employee: mason, carpenter, electrician	29	0
		self-employed worker: foreman, carpenter	6	0
	textile and craft industries	informal sector employee: dressmaker, cabinetmaker, brickwork, quinoa food processing	15	12
		self-employed worker: dressmaker, baker, musician	2	2
tertiary	trade and restaurant industries	informal sector employee: grocer, ironmonger, caretaker, waiter	14	10
		self-employed worker: street fooding, clothing trade, restaurant	14	37
	car services and transportation	informal sector employee: mechanic, welder, minibus driver, truck driver, railway worker	10	0
		self-employed worker: mechanic, bodywork, welding, minibus or truck business	4	0
	domestic work	informal sector employee: daytime or fulltime servant	0	50
	administration	civil servant and assimilated workers: nursing, rural professor, policeman, administrative officer	16	0

Source: Vassas-Toral (2015c)

As illustrated below (Figs. B-2, 3), many people practise a temporary migration and a circulation between these different places and their community. These practices attest to the maintaining of personal links with the rural place of origin and a strong territorial anchoring, which facilitated the return of emigrated people to their communities during the period of quinoa expansion.

Fig. B-2. Trajectory of residential mobility of Calixto, 28 y old, native of San Juan de Rosario, with residence in that community in 2008.

Source: Vassas-Toral, interviews and livelihood stories, 2007-2008.

The trajectory of Calixto is typical of many young people who emigrated for a long time in Argentina and Chile, before returning in their community to produce quinoa. After 1 year in Uyuni to complete his military service, and 9 years as a factory worker in Salta, Buenos Aires and Calama (1996-2006), Calixto returned in 2006 to San Juan where he is now living from quinoa production and touristic activities.

Fig. B-3.
Trajectory of residential mobility of Benedicto, 64 y old, native from Palaya, with doble residence in Palaya and Tupiza in 2008.

Source: Vassas-Toral, interviews and livelihood stories, 2007-2008.

The migration trajectory of Benedicto is a complex one. He left alone the community of Palaya when he was 11 years old and, while going to the school, began working in Chile as an agricultural worker (*peon*) on the farm of a family member, and then as a miner. After completing his military service at the age of 17 in La Paz, he settled in Palaya for 9 years, cultivating his own lands. However the income of agriculture was insufficient for the family to live. So Benedicto decided to leave his community and began a long migratory trajectory as school profesor in urban centers: Sucre, Tupiza, Iquique, Tarija, and again Tupiza. In 2008 Benedicto retired. After 24 years of absence, he returned in his native community to dedicate himself to quinoa production on the lands of which he had retained usufruct. Like numerous quinoa growers in the southern Altiplano, he practices now a circular mobility with doble residence between the countryside and the city.

Fig. B-4. Trajectory of circular mobility linked with the agricultural activity of Felix, 28 y old, native of San Juan de Rosario with residence in that community in 2008. Source: Vassas-Toral, interviews and livelihood stories, 2007-2008.

The trajectory of Felix is representative of numerous young people who contributed to the process of appropriation of lands in the southern altiplano. When he was 18 years old, Felix moved from his native community to Uyuni, then to Chile (Calama and Antofagasta) to work in the building sector. During eight years, he returned to his community every year to visit his parents. The link maintained with the community allowed him to move back in San Juan in 2006. However, having no direct access to cropland, Felix cultivated at first in sharecropping (*al partir*), before being able to clear some plots in the collective pasture lands, gradually becoming usufructuary of five hectares.

Fig. B-5. Trajectories of circular mobility linked with the agricultural activity of Teodoro and Graciela, both 71 y old, native of Otuyo with residence in La Paz in 2008. Source: Vassas-Toral, interviews and livelihood stories, 2007-2008.

1036 In 2008, Teodoro and Graciela were simultaneously residents in La Paz (the capital of Bolivia)
and quinoa growers in their native community of Otuyo, more than 500 km away. Their trajectories
1038 show a continuous circular mobility between the city and the community during more than twenty
years of migration (1963-1999). During this period, they didn't cultivate their lands any more but they
1040 returned once a year to their community to participate in local festivities, thus maintaining the right of
usufruct of their lands. From 1999 on, when Teodoro and Graciela decided to cultivate their lands
1042 again to produce quinoa for trade, the frequency of their circular mobility increased. They return three
or four times a year to cultivate their fields. The case of Teodoro and Graciela illustrates the
1044 emergence of "urban farmers" as a significant social group in the southern altiplano.

1046 **Cited references**

- Vassas-Toral, A., (2011) Ruralité et agriculture au prisme des mobilités spatiales. L'Altiplano Sud de
1048 Bolivie à l'heure du boom de la quinoa. PhD thesis. Université Paul Valéry - Montpellier III,
Montpellier, France, 345 p. + annexes. (in french)
- Vassas-Toral, A., (2015a) Movilidades de los productores de quinua y dinámicas territoriales en el
1050 Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), Racionalidades campesinas en
1052 los Andes del Sur: reflexiones en torno al cultivo de la quinua y otros vegetales andinos.
EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 231-280.
- Vassas-Toral, A., (2015b) Movilidades, recursos de tierra y sistema de producción agrícola en el
1054 Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), Racionalidades campesinas en
1056 los Andes del Sur: reflexiones en torno al cultivo de la quinua y otros vegetales andinos.
EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 281-321.
- Vassas-Toral, A. (2015c) Partir et cultiver: essor de la quinoa, mobilités et recompositions rurales en
1058 Bolivie. IRD Éditions, Marseille, France.
- 1060

Supplementary Material C. Localisation and cropland area in six communities of the southern
altiplano of Bolivia from 1963 to the 2000's.

Latitude and longitude are in decimal degrees. Cropland areas were calculated using aerial
photographs (1963), EROS satellite images (1972), and SPOT satellite images (2005-2008).

Community	Altitude (m)	Year		
		1963	1972	2005-2008 ¹
Chacoma	3700	0.0	0.0	0.0
	3750	20.1	46.6	450.0
	latitude S	3800	21.1	43.0
	-19.71	3850	54.4	107.0
	longitude W	3900	69.3	79.5
	-68.49	3950	109.0	124.1
		4000	31.8	27.6
		4050	26.0	21.6
Candelaria de Viluyo		4100	25.4	27.7
		4150	0.0	5.0
	3700	21.5	108.8	870.1
	3750	50.2	123.6	540.8
	3800	7.0	41.8	104.4
	latitude S	3850	0.6	3.2
	-19.68	3900	0.0	0.0
	longitude W	3950	0.0	0.0
Capura		4000	0.0	0.0
	-67.16	4050	0.0	0.0
		4100	0.0	0.0
	3700	0.0	0.0	0.0
	3750	1.0	2.5	15.1
	latitude S	3800	14.9	40.7
	-19.49	3850	2.6	4.6
	longitude W	3900	0.0	1.8
Chilalo		3950	0.0	0.0
	-67.28	3700	9.7	65.4
		3750	16.0	14.4
	latitude S	3800	17.5	10.5
	-19.8	3850	9.6	9.8
	longitude W	3900	16.0	5.2
	-67.81	3950	12.4	6.2
		4000	10.0	4.9
		4050	5.2	1.8
		4100	3.8	0.0
		4150	9.2	1.2
		4200	1.6	0.7

(continued)

¹ 2006 in Chacoma and Palaya, 2008 in Candelaria de Viluyo, 2005 in
Capura, Chilalo, and Otuyo.

(continued)

	Altitude (m)	1963	1972	2005-2008 ¹
Otuyo	3700	44.5	23.8	159.1
	3750	35.6	94.8	328.5
latitude S	3800	23.7	24.4	11.4
-19.59	3850	16.2	9.2	0.0
longitude W	3900	8.0	1.0	0.0
-67.76	3950	2.2	0.0	0.0
	4000	0.0	0.0	0.0
Palaya	3700	14.2	242.9	713.8
	3750	130.8	276.3	789.3
latitude S	3800	38.1	104.2	293.7
-19.48	3850	29.8	66.5	197.2
longitude W	3900	60.0	90.4	218.4
-68.08	3950	72.4	112.4	197.6
	4000	106.2	76.8	191.2
	4050	112.0	83.5	214.9
	4100	89.2	63.5	112.1
	4150	81.6	81.5	84.3
	4200	100.2	158.7	108.2
	4250	101.3	103.6	90.6
	4300	61.3	99.1	30.5
	4350	43.8	77.7	12.2
	4400	18.4	46.7	6.4
	4450	7.1	37.8	4.8
	4500	2.6	76.5	6.1

Supplementary Material D. The Vulnerability Scoping Diagram

The socio-environmental assessment of quinoa production in the southern Altiplano of Bolivia may have different formulations, depending on the structural unit and the process of change under assessment. The Vulnerability Scoping Diagram in fig. D-1 synthesizes such an assessment focused on the rural community as the exposure unit, and on ecological degradation as the possible source of unsustainability for the considered community (an alternative object of assessment could have been, for example, the farming unit exposed to the market dynamics). The diagram specifies the components of exposure, sensitivity and adaptive capacity identified in the system. In the present case, the agrosystem is exposed to real or potential risks related to climate imprevisibility, land degradation, as well as increased social inequity or the possible desaffection of part of the quinoa consumers. The sensitivity to these risks refers to agrotechnical components, such as the management of natural resources and the crop yield. It depends also on social and economic components like land access or the dependence towards actors external to the community.

In this context, the members of the community can show their adaptive capacity in different components: innovating in their individual and/or collective practices of land management, diversifying their markets, realizing pluri-activity with- or without temporary migration. The measures of these components can bring sustainability indicators. Examples of such indicators are: quantifying the expansion of the quinoa crop in the territory of the considered community, or characterizing the spatial and temporal dynamics of the mobility of the community members (Vassas-Toral, 2015a, b), or mapping frost risks at a geographical scale compatible with decision making for local crop management (Pouteau et al., 2011).

Fig. D-1. Vulnerability scoping diagram assessing the vulnerability to ecological degradation in the local quinoa production system.

1094 **Cited references**

- 1096 Pouteau, R., Rambal, S., Ratte, J.P., Gogé, F., Joffre, R., Winkel, T. (2011) Downscaling MODIS-
derived maps using GIS and boosted regression trees: the case of frost occurrence over the arid
Andean highlands of Bolivia. *Remote Sensing of Environment* 115, 117-129.
- 1098 Vassas-Toral, A., (2015a) Movilidades de los productores de quinua y dinámicas territoriales en el
Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), *Racionalidades campesinas en*
1100 *los Andes del Sur: reflexiones en torno al cultivo de la quinua y otros vegetales andinos.*
EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 231-280.
- 1102 Vassas-Toral, A., (2015b) Movilidades, recursos de tierra y sistema de producción agrícola en el
Altiplano Sur de Bolivia, in: Cruz, P., Joffre, R., Thierry, W. (Eds.), *Racionalidades campesinas en*
1104 *los Andes del Sur: reflexiones en torno al cultivo de la quinua y otros vegetales andinos.*
EdiUNJU, Universidad Nacional de Jujuy, Jujuy, Argentina, pp. 281-321.
- 1106
- 1108