



Unexpected high densities of the hybrid coral *Acropora prolifera* (Lamarck 1816) in Guadeloupe Island, Lesser Antilles

A Japaud, C Fauvelot, C Bouchon

► To cite this version:

A Japaud, C Fauvelot, C Bouchon. Unexpected high densities of the hybrid coral *Acropora prolifera* (Lamarck 1816) in Guadeloupe Island, Lesser Antilles. *Coral Reefs*, 2014, 33, pp.593 - 593. 10.1007/s00338-014-1169-7 . ird-03044174

HAL Id: ird-03044174

<https://ird.hal.science/ird-03044174>

Submitted on 7 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reef sites

Unexpected high densities of the hybrid coral *Acropora prolifera* (Lamarck 1816) in Guadeloupe Island, Lesser Antilles



Fig. 1 *Acropora prolifera* field in the Bay of Grand Cul-de-Sac Marin (a) and single *A. prolifera* colony (b)

Since the 1980s, coral populations of *Acropora palmata* (Lamarck 1816) and *A. cervicornis* (Lamarck 1816) have dramatically declined in the Caribbean and have been consequently classified as critically endangered since 2008 (IUCN).

Acropora palmata and *A. cervicornis* can form a viable F1 offspring previously described as *A. prolifera* by Lamarck (Vollmer and Palumbi 2002). Few isolated colonies of that morphotype have punctually been reported on Lesser Antilles reefs in the previous decades. In January 2013, high densities of morphologically identified *A. prolifera* colonies were observed in the Bay of Grand Cul-de-Sac Marin ($16^{\circ}21'24.19''\text{N}$; $61^{\circ}35'35.12''\text{W}$, Guadeloupe Island, Lesser Antilles), whereas no colonies were found at this exact same site prospected in May 2011 (Fig. 1). Genetic microsatellite analyses (unpublished data) verified the hybrid nature of the colonies and suggest that they may constitute a single clone. However, in other Caribbean sites, Fogarty (2010) found high genotypic diversity for that hybrid.

As a consequence of decreased parental species' density, eggs likely float unfertilized, increasing the probability of heterospecific sperm encounters and thus hybrid embryo formation (Fogarty et al. 2012).

References

- Fogarty ND (2010) Reproductive isolation and hybridization dynamics in threatened Caribbean acroporid corals. Ph.D. thesis, Florida State University, Tallahassee, Florida
- Fogarty ND, Vollmer SV, Levitan DR (2012) Weak prezygotic isolating mechanisms in threatened Caribbean *Acropora* corals. PLoS ONE 7:e30486
- Vollmer SV, Palumbi SR (2002) Hybridization and the evolution of reef coral diversity. Science 296:2023–2025

A. Japaud · C. Bouchon (✉)

UMR BOREA, CNRS 7208 – MNHN – UPMC – UCBN – IRD 207, Laboratoire d’Excellence « CORAIL », Université des Antilles et de la Guyane, BP 592, Pointe-à-Pitre, Guadeloupe, France
e-mail: claude.bouchon@univ-ag.fr

C. Fauvelot

UR 227 CoReUs 2, Laboratoire d’Excellence « CORAIL », Institut de Recherche pour le Développement (IRD), Noumea, New Caledonia

Received: 26 March 2014 / Accepted: 14 May 2014
© Springer-Verlag Berlin Heidelberg 2014

Coral Reefs (2014)
DOI 10.1007/s00338-014-1169-7